



## ABSTRACTS OF TALKS

---

**Mirosław Adamek**  
University of Bielsko-Biala, Poland

### On Hermite-Hadamard type inequalities for $F$ -convex functions

Let  $I$  be a nonempty and open interval of  $\mathbb{R}$  and  $F: \mathbb{R} \rightarrow \mathbb{R}$  be a fixed function. A function  $f: I \rightarrow \mathbb{R}$  will be called  $F$ -convex if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) - t(1-t)F(x-y)$$

for all  $x, y \in I$  and  $t \in (0, 1)$ .

The classical Hermite-Hadamard inequality says that for any convex function  $f: I \rightarrow \mathbb{R}$  we have

$$f\left(\frac{x+y}{2}\right) \leq \frac{1}{y-x} \int_x^y f(u)du \leq \frac{f(x)+f(y)}{2}$$

for all different  $x, y \in I$ .

In this talk we present Hermite-Hadamard type inequalities for  $F$ -convex functions.

As a consequence of our investigations we get the following inequality

$$f\left(\frac{x+y}{2}\right) + \frac{1}{y-x} \int_x^y F\left(u - \frac{x+y}{2}\right) du \leq \frac{1}{y-x} \int_x^y f(u)du \leq \frac{f(x)+f(y)}{2} - \frac{1}{6}F(x-y)$$

for all different  $x, y \in I$ .

**Nutefe Kwami Agbeko**  
University of Miskolc, Hungary

## On some lattice-valued equations and inequalities

Since early 90's we have considered lattice-valued functions and operators defined on diverse sets with algebraic structures, by replacing the addition with lattice operations. In this perspective, the addition in the definition of the probability measure as well as in the definition of the Lebesgue's integral (or mathematical expectation) is substituted with the supremum operation and the so-defined functions (*optimal measure*, resp. *optimal average*) behave similarly like their counterparts in Measure Theory. From 2012 the Cauchy functional equation has also been studied for lattice-valued functions (which we can term *lattice-valued Cauchy functional equation*), where the supremum replaces the addition in the Cauchy functional equation, focusing on the Ulam's type stability. The main goal of our presentation is to talk about morphisms between sets with an algebraic structure and an order structure, through associated functional equations and inequalities: we discuss the separation problem for the inequalities and the Hyers-Ulam stability of the main equation.

### References

- [1] N.K. Agbeko, *On optimal averages*, Acta Math. Hung. **63** (1-2) (1994), 1–15.
- [2] N.K. Agbeko, *On the structure of optimal measures and some of its applications*, Publ. Math. Debrecen **46** (1-2) (1995), 79–87.
- [3] N.K. Agbeko, *Stability of maximum preserving functional equations on Banach lattices*, Miskolc Math. Notes, **13** No. 2 (2012), 187–196.
- [4] N.K. Agbeko and S.S. Dragomir, *The extension of some Orlicz space results to the theory of optimal measure*, Math. Nachr. **286** No 8-9 (2013), 760–771.
- [5] N.K. Agbeko, *The Hyers-Ulam-Aoki type stability of some functional equation on Banach lattices*, Bull. Polish Acad. Sci. Math. **63** No. 2 (2015), 177–184.
- [6] N.K. Agbeko, *A remark on a result of Schwaiger*, Indag. Math. **28**, Issue 2, (2017), 268–275.
- [7] N.K. Agbeko, W. Fechner and E. Rak, *On lattice-valued maps stemming from the notion of optimal average*. Acta Math. Hungar. **152** (2017), 72–83.

**Pekka Alestalo**  
Aalto University, Finland

## Sharp extension results for bilipschitz maps

An  $L$ -bilipschitz map  $f: A \rightarrow \mathbf{R}^n$  satisfies the double inequality

$$\|x - y\|/L \leq \|f(x) - f(y)\| \leq L\|x - y\|$$

for all  $x, y \in A \subset \mathbf{R}^n$ . For topological reasons, these maps cannot usually be extended to a homeomorphism  $F: \mathbf{R}^n \rightarrow \mathbf{R}^n$ . In the other extreme, it is sometimes possible to find an  $L$ -bilipschitz extension  $F$  with the same constant  $L$ . We present some positive results and examples related to this problem.

### References

- [1] P. Alestalo, D.A. Trotsenko, *Radial extensions of bilipschitz maps between unit spheres*, Siberian Electronic Mathematical Reports 15, 839–843 (2018).  
DOI 10.17377/semi.2018.15.071

**Alina Ramona Baias**  
Technical University of Cluj-Napoca, Romania

## On the best Ulam constant of a third order linear difference equation

Let  $X$  be a Banach space over the field  $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ . We give a result on Ulam stability for the linear difference equation

$$x_{n+3} = ax_{n+2} + bx_{n+1} + cx_n, \quad n \geq 0, \tag{1}$$

where  $a, b, c \in \mathbb{K}$ ,  $x_0, x_1, x_2 \in X$ . Moreover, if all the roots of the characteristic equation of (1) have the modulus greater than 1, we obtain the best Ulam constant of the equation.

### References

- [1] J. Brzdek, D. Popa, I. Raşa, B. Xu, *Ulam Stability of Operators*, Academic Press, (2018).  
[2] A.R. Baias, F. Blaga, D. Popa, *Best Ulam constant for a linear difference equation*, Carpathian J. Math. **35** (2019), 13–22.  
[3] A.R. Baias, D. Popa, *On Ulam stability of a linear difference equation in Banach spaces*, Bull. Malays. Math. Sci. Soc., DOI: 10.1007/s40840-019-00744-6.  
[4] J. Brzdek, D. Popa, I. Raşa, *Hyers Ulam stability with respect to gauges*, J. Math. Anal. Appl. **453** (2017), 620–628, .

**Karol Baron**  
Uniwersytet Śląski, Poland

## Remarks on continuous solutions of an iterative functional equation

Assume  $X$  is a real separable Hilbert space,  $\Lambda : X \rightarrow X$  is linear and continuous with  $\|\Lambda\| < 1$ , and  $\mu$  is a probability Borel measure on  $X$  with finite first moment. We examine continuous at zero solutions  $\varphi : X \rightarrow \mathbb{C}$  of the equation

$$\varphi(x) = \hat{\mu}(x)\varphi(\Lambda x).$$

**Liviu Cădariu-Brăiloiu**  
Politehnica University of Timișoara, Romania

## Generalized Hyers-Ulam stability of some functional equations

The aim of this talk is to present several generalized Hyers-Ulam stability properties for some functional equations, by using the fixed point method.

### References

- [1] J. Brzdęk, L. Cădariu, K. Ciepliński, *Fixed point theory and the Ulam stability*, J. Function Spaces **2014** (2014), Article ID 829419, 16 pp.
- [2] J. Brzdęk, L. Cădariu, *Stability for a family of equations generalizing the equation of  $p$ -Wright affine functions*, Applied Mathematics and Computation **276** (2016), 158–171.
- [3] K. Ciepliński, *Applications of fixed point theorems to the Hyers-Ulam stability of functional equations – a survey*, Ann. Funct. Anal. **3**(1) (2012), 151–164.

# Jacek Chudziak

University of Rzeszów, Poland

## Convexity and quasi-convexity of the zero utility principle

Assume that  $\mathcal{X}_+$  is a family of all non-negative bounded random variables on a given probability space. The elements of  $\mathcal{X}_+$  represent the risks to be insured by an insurance company. An important question of the theory of insurance risk premiums is to assign to every  $X \in \mathcal{X}_+$  a premium for the insurance contract. One of the methods of determining the premium is *the zero utility principle*. It defines the premium for a risk  $X \in \mathcal{X}_+$  as a real number  $H_u(X)$  satisfying equation

$$E[u(H_u(X) - X)] = 0 \tag{1}$$

where  $u : \mathbb{R} \rightarrow \mathbb{R}$  is a strictly increasing continuous function with  $u(0) = 0$ . It turns out that, for every  $X \in \mathcal{X}_+$ , such a number  $H_u(X)$  exists and it is unique. Therefore equation (1) determines in an implicit way a functional  $H_u : \mathcal{X}_+ \rightarrow \mathbb{R}$ .

Several results concerning properties of the functional  $H_u$  can be found e.g. in [1, 2, 3] and [6]. It is known that, for every strictly increasing continuous function  $u : \mathbb{R} \rightarrow \mathbb{R}$  satisfying  $u(0) = 0$ ,  $H_u$  is monotone and conditionally translation invariant. However, in general, it is not convex. In this talk we present a characterization of convexity and quasi-convexity of  $H_u$ . A fundamental role in our investigations is played by *quasideviation means* (cf. [4, 5]).

### References

- [1] N. L. Bowers, H. U. Gerber, J. C. Hickman, D. A. Jones and C. J. Nesbitt, *Actuarial Mathematics*, The Society of Actuaries, Itasca, Illinois, 1986.
- [2] H. Bühlmann, *Mathematical Models in Risk Theory*, Springer, Berlin, 1970.
- [3] H. U. Gerber, *An Introduction to Mathematical Risk Theory*, S. S. Huebner Foundation, R. D. Irwin Inc., Homewood Illinois, 1979.
- [4] Zs. Páles, *Characterization of quasideviation means*, Acta. Math. Sci. Hungar. **40** (1982), 243–260.
- [5] Zs. Páles, *General inequalities for quasideviation means*, Aequationes Math. **36** (1988), 32–56.
- [6] T. Rolski, H. Schmidli, V. Schmidt and J. Teugels, *Stochastic Processes for Insurance and Finance*, John Wiley & Sons, New York 1999.

# Włodzimierz Fechner

Lodz University of Technology, Poland

## Sincov's inequalities on topological spaces

During the talk we will discuss the multiplicative Sincov's inequality:

$$G(a, b) \leq G(a, x) \cdot G(x, b), \quad a, x, b \in X.$$

We assume that  $X$  is a topological space and  $G$  is a continuous map. We also study the reverse inequality:

$$F(a, b) \geq F(a, x) \cdot F(x, b), \quad a, x, b \in X$$

and the additive version of the original inequality:

$$H(x, z) \leq H(x, y) + H(y, z), \quad x, y, z \in X.$$

A corollary for generalized metric is derived.

### References

- [1] Włodzimierz Fechner, *Sincov's inequalities on topological spaces* (manuscript), arXiv: 1811.00303v3 [math.FA] 21 Nov 2018, 10 pp.

## Roman Ger

Silesian University of Katowice, Poland

### On convex type functional inequalities

The lecture focuses on some weak regularity requirements forcing the automatic continuity of real convex functionals on normed spaces and, more generally, on locally convex linear topological spaces, supporting and separation theorems (i.e. geometric counterparts of various generalizations of the celebrated Hahn-Banach extension theorem). In some situations, groups will be considered as potential domains and the related problem of their classification with respect to the question whether or not they admit invariant means (amenability).

Because of the significant role and good geometry of strictly convex spaces some characterizations of them in terms of solutions of the fundamental Cauchy functional equation assumed to be satisfied *modulo* a norm with the emphasis given to the factorization of these solutions into the additive and isometric parts. Several further necessary and sufficient conditions for the strict convexity of the given space will be analyzed as well along with some characterizations of inner product spaces with the aid of suitable iterations of the difference operators.

Delta-convex mappings between normed linear spaces provide a generalization of functions which are representable as a difference of two convex functions to the case of vector valued maps. Following L. Veselý and L. Zajíček, (*Delta-convex mappings between Banach spaces and applications*, Dissertationes Math. 289, Polish Scientific Publishers, Warszawa, 1989) we show that this class of mappings has very good properties proving that a generalization proposed is well established. Strict connections with the Hyers-Ulam stability in the theory of functional equations and inequalities will be revealed. We look also for possibly mild regularity conditions upon the maps whose vector convex differences are controlled by their scalar counterparts, forcing these maps to be delta-convex. Finally, vector analogues of the celebrated Hermite-Hadamard type inequalities will also be presented.

## Moshe Goldberg

Technion-Israel Institute of Technology, Israel

### Extending the spectral radius to finite-dimensional power-associative algebras

The purpose of this talk is to introduce a new concept, the *radius* of elements in arbitrary finite-dimensional power-associative algebras over the field of real or complex numbers. It is an extension of the well known notion of the spectral radius.

As examples, we shall discuss this new radius in the setting of matrix algebras, where it indeed reduces to the spectral radius, and then in the Cayley-Dickson algebras, where it is something quite different.

# Richárd Grünwald

University of Debrecen, Hungary

## Characterization of the equality of generalized Bajraktarević means

The purpose of the talk is to investigate the equality problem of generalized Bajraktarević means, i.e., to solve the functional equation

$$f^{(-1)}\left(\frac{p_1(x_1)f(x_1) + \cdots + p_n(x_n)f(x_n)}{p_1(x_1) + \cdots + p_n(x_n)}\right) = g^{(-1)}\left(\frac{q_1(x_1)g(x_1) + \cdots + q_n(x_n)g(x_n)}{q_1(x_1) + \cdots + q_n(x_n)}\right), \quad (*)$$

which holds for all  $x = (x_1, \dots, x_n) \in I^n$ , where  $n \geq 2$ ,  $I$  is a nonempty open real interval, the unknown functions  $f, g : I \rightarrow \mathbb{R}$  are strictly monotone,  $f^{(-1)}$  and  $g^{(-1)}$  denote their generalized left inverses, respectively, and  $p = (p_1, \dots, p_n) : I \rightarrow \mathbb{R}_+^n$  and  $q = (q_1, \dots, q_n) : I \rightarrow \mathbb{R}_+^n$  are also unknown functions. This equality problem in the symmetric two-variable (i.e., when  $n = 2$ ) case was already investigated and solved under sixth-order regularity assumptions by Losonczi in 1999. In the nonsymmetric two-variable case, assuming three times differentiability of  $f, g$  and the existence of  $i \in \{1, 2\}$  such that either  $p_i$  is twice continuously differentiable and  $p_{3-i}$  is continuous on  $I$ , or  $p_i$  is twice differentiable and  $p_{3-i}$  is once differentiable on  $I$ , we prove that (\*) holds if and only if there exist four constants  $a, b, c, d \in \mathbb{R}$  with  $ad \neq bc$  such that

$$cf + d > 0, \quad g = \frac{af + b}{cf + d}, \quad \text{and} \quad q_\ell = (cf + d)p_\ell \quad (\ell \in \{1, \dots, n\}). \quad (1)$$

In the case  $n \geq 3$ , we obtain the same conclusion with weaker regularity assumptions. Namely, we suppose that  $f$  and  $g$  are three times differentiable,  $p$  is continuous and there exist  $i, j, k \in \{1, \dots, n\}$  with  $i \neq j \neq k \neq i$  such that  $p_i, p_j, p_k$  are differentiable.

### References

- [1] J. Aczél and Z. Daróczy, *Über verallgemeinerte quasilineare Mittelwerte, die mit Gewichtsfunktionen gebildet sind*, Publ. Math. Debrecen **10** (1963), 171–190.
- [2] M. Bajraktarević, *Sur une équation fonctionnelle aux valeurs moyennes*, Glasnik Mat.-Fiz. Astronom. Društvo Mat. Fiz. Hrvatske Ser. II **13** (1958), 243–248.
- [3] M. Bajraktarević, *Sur une généralisation des moyennes quasilineaires*, Publ. Inst. Math. (Beograd) (N.S.) **3** (17) (1963), 69–76.
- [4] Z. Daróczy and L. Losonczi, *Über den Vergleich von Mittelwerten*, Publ. Math. Debrecen **17** (1970), 289–297.
- [5] Z. Daróczy and Zs. Páles, *On comparison of mean values*, Publ. Math. Debrecen **29** (1982), 107–115.
- [6] C. Gini, *Di una formula compressiva delle medie*, Metron **13** (1938), 3–22.
- [7] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge University Press, Cambridge, 1934, (first edition), 1952 (second edition).
- [8] L. Losonczi, *Equality of two variable weighted means: reduction to differential equations*, Aequationes Math. **58** (1999) 223–241.
- [9] L. Losonczi, *Equality of two variable means revisited*, Aequationes Math. **71** (2006), 228–245.
- [10] L. Losonczi, *Homogeneous non-symmetric means of two variables*, Demonstratio Math. **40** (2007), 169–180.
- [11] L. Losonczi, *Homogeneous symmetric means of two variables*, Aequationes Math. **74** (2007), 262–281.
- [12] Zs. Páles and A. Zakaria, *On the equality of Bajraktarević means to quasi-arithmetic means*, (2019), submitted.



## László Horváth

University of Pannonia, Hungary

### Sharp Gronwall-Bellman type integral inequalities with delay

Various attempts have been made to give an upper bound for the solutions of the delayed version of the Gronwall-Bellman integral inequality, but the obtained estimations are not sharp. In this talk a new approach is presented to get sharp estimations for the nonnegative solutions of the considered delayed inequalities. The results are based on the idea of the generalized characteristic inequality. Our method gives sharp estimation, and therefore the results are more exact than the earlier ones.

## Eliza Jabłońska

Pedagogical University of Kraków, Poland

### Haar 'small' sets in abelian Polish groups

It is well known [1] that a subset  $A$  of an abelian Polish group  $X$  is called *Haar null* if there are a universally measurable set  $B \subset X$  with  $A \subset B$  and a Borel probability measure  $\mu$  on  $X$  such that  $\mu(x + B) = 0$  for all  $x \in X$ . In [2] Darji introduced another family of "small" sets in an abelian Polish group  $X$ ; he called a set  $A \subset X$  *Haar meager* if there is a Borel set  $B \subset X$  with  $A \subset B$ , a compact metric space  $K$  and a continuous function  $f : K \rightarrow X$  such that

$$f^{-1}(B + x) \text{ is meager in } K \text{ for every } x \in X.$$

In a locally compact group these two definitions are equivalent to definitions of Haar measure zero sets and meager sets, respectively. That is why we can say that the notion of a Haar meager set is a topological analog to the notion of a Haar null set. Since lots of similarities between meager sets and sets of Haar measure zero are well known in locally compact groups (see e.g. [3]), we would like to present some analogies between Haar meager sets and Haar null sets.

### References

- [1] J.P.R. Christensen, *On sets of Haar measure zero in abelian Polish groups*, Israel J. Math. **13** (1972), 255–260.
- [2] U.B. Darji, *On Haar meager sets*, Topology Appl. **160** (2013), 2396–2400.
- [3] J.C. Oxtoby, *Measure and Category*, Springer-Verlag, New York-Heidelberg-Berlin, 1971.

## Suresh Kumar Murugan

The Gandhigram Rural Institute-Deemed University, India

### Series-Like Iterative Functional Equation for PM Functions

It has been proved that the series-like iterative functional equation of the form  $\sum_{n=1}^{\infty} \lambda_n f^n(x) = F(x)$  has solution provided  $F$  is strictly monotone. In this paper, we extend this problem for the class of piecewise monotone functions. We obtained sufficient conditions on the existence of continuous solutions for the class of piecewise monotone functions using the method of characteristic interval. Also as an application we construct a everywhere continuous nowhere differentiable function on the real line.

#### References

- [1] V. Murugan and P. V. Subrahmanyam. Existence of solutions for equations involving iterated functional series. *Fixed Point Theory Appl.* 2, 219–232, 2005.
- [2] V. Murugan and P. V. Subrahmanyam. Existence of continuous solutions for an iterative functional series equation with variable coefficients. *Aequations Math.*, 78, 167-176, 2009.
- [3] L. Liu, X. Gong. The polynomial-like iterative equation for PM functions. *Sci. China Math.* 60 (8), 1503–1514, 2017.
- [4] Y.J Cho, M. Suresh Kumar, and V. Murugan. Iterative Roots of Non-PM Functions and Denseness. *Results Math* 73:13, 2018.

## Zbigniew Leśniak

Pedagogical University of Kraków, Poland

### On fixed points of functions with values in a dq-metric space

(joint work with JANUSZ BRZDĘK AND EL-SAYED EL-HADY)

We present a fixed-point theorem for an operator acting on some classes of functions with values in a dq-metric space and show its applications to prove the stability in Ulam sense of some types of functional and difference equations. The fixed points of such operators turn out to be exact solutions of the considered equations that meet the imposed conditions.

#### References

- [1] J. Brzdęk, El-s. El-hady, Z. Leśniak, *On fixed points of a linear operator of polynomial form of order 3*, J. Fixed Point Theory Appl. 20 (2018), Article:85, 10 pp.
- [2] J. Brzdęk, El-s. El-hady, Z. Leśniak, *On fixed-point theorem in classes of function with values in a dq-metric space*, J. Fixed Point Theory Appl. 20 (2018), Article:143, 16 pp.

## Renata Malejki

Pedagogical University of Kraków, Poland

### On Ulam stability of a generalization of the Fréchet functional equation on a restricted domain

In this paper we prove the Ulam type stability of a generalization of the Fréchet functional equation on a restricted domain. In the proofs the main tool is a fixed point theorem for some function spaces.

#### References

- [1] J. Brzdęk, Z. Leśniak, R. Malejki, *On the generalized Fréchet functional equation with constant coefficients and its stability*, *Aequationes Math.* **92** (2018), 355–373 .
- [2] R. Malejki, *Stability of a generalization of the Fréchet functional equation*, *Ann. Univ. Paedagog. Crac. Stud. Math.* **14** (2015), 69–79 .

## Janusz Matkowski

University of Zielona Góra, Poland

### Quasi-Cauchy difference means

Quasi-Cauchy difference means of:

(i) additive type, i.e. the functions of the form

$$M_f(x_1, \dots, x_k) = F^{-1}(f(x_1 + \dots + x_k) - (f(x_1) + \dots + f(x_k))),$$

where  $F(x) = f(kx) - kf(x)$ ;

(ii) exponential type, i.e. the functions of the form

$$E_f(x_1, \dots, x_k) = F^{-1}\left(f\left(\sum_{j=1}^k x_j\right) - \prod_{j=1}^k f(x_j)\right),$$

where  $F(x) = f(kx) - [f(x)]^k$ ;

(iii) logarithmic type, i.e. the functions of the form

$$L_f(x_1, \dots, x_k) = F^{-1}\left(f\left(\prod_{j=1}^k x_j\right) - \sum_{j=1}^k f(x_j)\right)$$

where  $F(x) := f(x^k) - kf(x)$ ;

(iv) of power type, i.e. the functions of the form

$$P_f(x_1, \dots, x_k) = F^{-1}\left(f\left(\prod_{j=1}^k x_j\right) - \prod_{j=1}^k f(x_j)\right),$$

where  $F(x) := f(x^k) - [f(x)]^k$ , as well as the respective functions with difference replaced by division, will be considered.

## Janusz Morawiec

University of Silesia, Poland

### Around a Kazimierz Nikodem result - part I

(joint work with THOMAS ZÜRCHER)

Let  $(X, \mathcal{A}, \mu)$  be a probability space and let  $S: X \rightarrow X$  be a measurable transformation. Motivated by the paper of K. Nikodem [1], we concentrate on a functional equation generating measures that are absolutely continuous with respect to  $\mu$  and  $\varepsilon$ -invariant under  $S$ . As a consequence of the investigation, we obtain a result on the existence and uniqueness of solutions  $\varphi \in L^1([0, 1])$  of the functional equation

$$\varphi(x) = \sum_{n=1}^N |f'_n(x)| \varphi(f_n(x)) + g(x),$$

where  $g \in L^1([0, 1])$  and  $f_1, \dots, f_N: [0, 1] \rightarrow [0, 1]$  are functions satisfying some extra conditions. The results we are going to present were recently published in [2].

#### References

- [1] K. Nikodem, *On  $\varepsilon$ -invariant measures and a functional equation*, Czechoslovak Math. J. **41**(116) (4) (1991) 565–569.
- [2] J. Morawiec and T. Zürcher, *An application of functional equations for generating  $\varepsilon$ -invariant measures*, J. Math. Anal. Appl. **476** (2019), 759-772.

## Jacek Mrowiec

University of Bielsko-Biała, Poland

### On strongly convex functions of higher order

Two given finite sequences  $(b_1, \dots, b_m)$  and  $(c_1, \dots, c_m)$ , where  $m \in \mathbb{N}$ ,  $b_k \in \{0, 1\}$ ,  $c_k > 0$ ,  $k = 1, \dots, m$ , describe a function  $F$  defined on a bounded interval  $I$  in the following way:

$b_k = 1$  means that  $F$  is strongly convex of order  $k$  with modulus  $c_k$  on  $I$ ,

$b_k = 0$  means that  $F$  is strongly concave of order  $k$  with modulus  $c_k$  on  $I$ ,  $k = 1, \dots, m$ .

For any fixed pair of such sequences the question of existence of  $F$  arises. In the talk the construction of the example of a function with the desired property will be presented. The case of infinite sequences also will be considered.

**Kazimierz Nikodem**  
University of Bielsko-Biala, Poland

## On strongly convex functions and related classes of functions

Let  $D$  be a convex subset of a normed space and  $c > 0$ . A function  $f : D \rightarrow \mathbb{R}$  is called *strongly convex* with modulus  $c$  if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) - ct(1-t)\|x-y\|^2$$

for all  $x, y \in D$  and  $t \in [0, 1]$ ;  $f$  is called *strongly midconvex* with modulus  $c$  if

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x)+f(y)}{2} - \frac{c}{4}\|x-y\|^2, \quad x, y \in D.$$

Strongly convex functions are useful in optimization theory and mathematical economics. Many properties and applications of them can be found in the literature. In my talk some results on strongly convex functions and related classes of functions obtained by the author with co-authors in the last few years are presented. In particular, discrete and integral Jensen-type inequalities and a Hermite–Hadamard–type inequality for strongly convex functions are obtained. Counterparts of the classical Bernstein–Doetsch and Sierpiński theorems for strongly midconvex functions are given. New characterizations of inner product spaces involving strong convexity are obtained. A representation of strongly Wright-convex functions and a characterization of functions generating strongly Schur-convex sums are presented. Finally, some properties of strongly convex set-valued maps and strongly convex stochastic processes are presented.

## Diana Otrocol

Technical University of Cluj-Napoca, Romania,  
T. Popoviciu Institute of Numerical Analysis, Romanian Academy, Romania

## Functional equations and entropies

We consider entropies corresponding to some probability distributions and establish functional/differential equations satisfied by them. Connections between these entropies are studied. As applications we investigate shape properties of the entropies and derive combinatorial identities.

## Lahbib Oubbi

Mohammed V University in Rabat, Ecole Normale Supérieure, Morocco

### Hyers-Ulam stability and hyperstability of a general functional equation in random normed spaces, a purely fixed point approach

If  $X$  is a real or complex vector space,  $(Y, F, T_M)$  is a Random normed space, and  $f : X \rightarrow Y$  a mapping, then using a purely fixed point approach, we prove the Ulam-Hyers stability and hyperstability of the general functional equation

$$\sum_{i=1}^m A_i f\left(\sum_{j=1}^n a_{ij} x_j\right) + A = 0.$$

Here  $f$  is a mapping from  $X$  into a Random normed space  $(Y, F, T_M)$ ,  $m$  and  $n$  are positive integers, for every  $i \in \{1, \dots, m\}$  and  $j \in \{1, \dots, n\}$ ,  $A_i$  and  $a_{ij}$  are scalars, and  $A$  is a vector from  $Y$ . Several known results can be derived.

#### References

- [1] A. Bahyrycz, J. Olko, *On stability of the general linear equation*, Aequat. Math. **89** (2015), 1461–1474.
- [2] A. Bahyrycz, J. Olko, *Hyperstability of general linear functional equation*, Aequat. Math. **90** (2016), 527–540.
- [3] J. Brzdęk, J. Chudziak, Zs. Páles, *A fixed point approach to stability of functional equations*, Nonlinear Anal. **74** (2011), 6728–6732.
- [4] Z. Dong, *On Hyperstability of Generalised Linear Functional Equations in Several Variables*. Bull. Aust. Math. Soc. **92** (2015), 259–267.

## Zsolt Páles

University of Debrecen, Hungary

### Optimal error functions for approximately monotone and convex functions

Let  $I$  be a nonempty open real interval and let  $\ell(I) \in ]0, \infty]$  denote its length. Given a nonnegative error function  $\Phi : [0, \ell(I)[ \rightarrow \mathbb{R}_+$ , a function  $f : I \rightarrow \mathbb{R}$  will be called a  $\Phi$ -monotone function if, for all  $x, y \in I$  with  $x \leq y$ ,

$$f(x) \leq f(y) + \Phi(y - x).$$

We say that a function  $f : I \rightarrow \mathbb{R}$  is  $\Phi$ -convex if, for all  $x, y \in I$  and  $t \in [0, 1]$ , it satisfies the inequality

$$f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y) + t\Phi((1 - t)|x - y|) + (1 - t)\Phi(t|x - y|).$$

In the talk, we discuss the following problem: If  $\Phi : [0, \ell(I)[ \rightarrow \mathbb{R}_+$  is an error function then determine the smallest error function  $\Phi^* : [0, \ell(I)[ \rightarrow \mathbb{R}_+$  such that  $\Phi$ -monotonicity and  $\Phi$ -convexity imply  $\Phi^*$ -monotonicity and  $\Phi^*$ -convexity, respectively.

## Rajendra Pant

University of Johannesburg, South Africa

### Viscosity Approximation methods for multi-valued nonexpansive mappings

We present some viscosity approximation theorems for multi-valued generalized nonexpansive mappings with applications to variational inequality, split common fixed point, split common null point and convex mathematical programming problems. Some numerical computations will be presented to illustrate our results.

## Paweł Pasteczka

Pedagogical University of Kraków, Poland

### Weakening of Hardy property for means

The aim of this talk is to find a broad family of means defined on a subinterval of  $I \subset [0, +\infty)$  such that

$$a_1 + \mathcal{M}(a_1, a_2) + \mathcal{M}(a_1, a_2, a_3) + \cdots < +\infty \quad \text{for all } a \in \ell_1(I).$$

Equivalently, the averaging operator

$$(a_1, a_2, a_3, \dots) \mapsto (a_1, \mathcal{M}(a_1, a_2), \mathcal{M}(a_1, a_2, a_3), \dots)$$

is a selfmapping of  $\ell_1(I)$ . This property is closely related to so-called Hardy inequality for means (which additionally requires boundedness of this operator).

We prove that these two properties are equivalent in a broad family of Gini means. Moreover, it is shown that this is not the case for quasi-arithmetic means. However, weak-Hardy property is localizable for this family.

## Dorian Popa

Technical University of Cluj-Napoca, Romania

### Ulam stability of an operatorial difference equation

Let  $T$  be a bounded linear operator acting on a Banach space  $X$ . We obtain some results on Ulam stability for the linear difference equation  $x_{n+1} = Tx_n + a_n$  associated to an iterative process for the linear equation  $x - Tx = y$ . As applications we get some stability results for the case when  $X$  is a finite dimensional space and for the case when  $T$  is Fredholm operator.

#### References

- [1] A.R. Baias, D. Popa, *On Ulam stability of a linear difference equation in Banach spaces*, Bull. Malays. Math. Sci. Soc., DOI: 10.1007/s40840-019-00744-6.
- [2] J. Brzdek, D. Popa, B. Xu, *On nonstability of the linear recurrence of order one*, J. Math. Anal. Appl. **367** (2010), 146–153.
- [3] J. Brzdek, D. Popa, I. Raşa, *Hyers Ulam stability with respect to gauges*, J. Math. Anal. Appl. **453** (2017), 620–628.
- [4] J. Brzdek, D. Popa, I. Raşa, B. Xu, *Ulam Stability of Operators*, Academic Press, (2018).
- [5] C. Buşe, D. O'Regan, O. Sailerli, A. Tabassum, *Hyers-Ulam stability and discrete dichotomy for difference periodic systems*, Bull. Sci. Math. **140** (2016), 908–934.



# Teresa Rajba

University of Bielsko-Biala, Poland

## On some inequalities for Bernstein operators and convex functions

For  $n \in \mathbb{N}$ , the **Bernstein basic polynomials** are given as follows

$$b_{n,i}(x) = \binom{n}{i} x^i (1-x)^{n-i} \quad (i = 0, 1, \dots, n, x \in [0, 1]),$$

the **classical Bernstein operators**  $B_n : \mathbb{C}([0, 1]) \rightarrow \mathbb{C}([0, 1])$ , are defined by

$$(B_n f)(x) = \sum_{i=0}^n b_{n,i}(x) f\left(\frac{i}{n}\right) \quad (x \in [0, 1]).$$

The following inequality was conjectured as an open problem by I. Raşa in [3]

$$\sum_{i,j=0}^n (b_{n,i}(x)b_{n,j}(x) + b_{n,i}(y)b_{n,j}(y) - 2b_{n,i}(x)b_{n,j}(y)) f\left(\frac{i+j}{2n}\right) \geq 0 \quad (1)$$

for each convex function  $f \in \mathbb{C}([0, 1])$  and for all  $x, y \in [0, 1]$ . The proof of inequality (1) was given in [2]. Raşa [4] remarked, that (1) is equivalent to

$$(B_{2n} f)(x) + (B_{2n} f)(y) \geq 2 \sum_{i=0}^n \sum_{j=0}^n b_{n,i}(x)b_{n,j}(y) f\left(\frac{i+j}{2n}\right). \quad (2)$$

In [1] we give some generalizations of inequality (2).

### References

- [1] A. Komisarski, T. Rajba, *Convex order for convolution polynomials of Borel measures*, arXiv: 1811.03827v1 [math.CA] 9 Nov 2018.
- [2] J. Mrowiec, T. Rajba, S. Waśowicz, *A solution to the problem of Raşa connected with Bernstein polynomials*, J. Math. Anal. Appl., **446** (2017), 864–878.
- [3] I. Raşa, 2. Problem, p. 164. In: Report of Meeting Conference on Ulam's Type Stability, Rytro, Poland, June 2–6, 2014, Ann. Univ. Paedagog. Crac. Stud. Math. **13** (2014), 139–169.
- [4] I. Raşa, *Bernstein polynomials and convexity: recent probabilistic and analytic proofs*, The Workshop "Numerical Analysis, Approximation and Modeling", T. Popoviciu Institute of Numerical Analysis, Cluj-Napoca, June 14, 2017, <http://ictp.acad.ro/zileleacademice-clujene-2017/>.

## Ioan Raşa

Technical University of Cluj-Napoca, Romania

### Functional equations and inequalities for the index of coincidence

Let  $(p_0(x), p_1(x), \dots)$  be a probability distribution depending on a real parameter  $x$ . The associated index of coincidence is  $S(x) := \sum_{k=0}^{\infty} (p_k(x))^2$ . The Rényi entropy and the Tsallis entropy are defined by  $R(x) := -\log S(x)$  and  $T(x) := 1 - S(x)$ . Starting with the binomial distribution, we establish functional equations and inequalities and use them to investigate convexity properties of the functions  $S(x)$ ,  $R(x)$  and  $T(x)$ . Applications and new open problems are mentioned.

#### References

- [1] I. Raşa, *Convexity properties of some entropies (II)*, Preprint 2019.

## Debmalya Sain

Indian Institute of Science, India

### Norm attainment set of a bounded linear operator between Banach spaces

It is a topic of current interest in the geometry of Banach spaces to study the norm attainment set of a bounded linear operator between Banach spaces. In this talk, I would like to explore the various facets of this problem, including the case of bounded linear operators between Hilbert spaces and Banach spaces. We would show that it is possible to completely characterize Euclidean spaces among Minkowski spaces, in terms of the operator norm attainment set. We would further explore the norm attainment set of a bounded linear operator between Banach spaces. Using the concept of Birkhoff-James orthogonality and semi-inner-products in Banach spaces, we completely characterize the operator norm attainment set in the setting of Banach spaces. If time permits, we would also like to briefly mention the various areas of application of the norm attainment set of a bounded linear operator, including the study of extreme contractions.

#### References

- [1] D. Sain, *On the norm attainment set of a bounded linear operator*, J. Math. Anal. Appl., **457** (2018), 67–76.

**Ekaterina Shulman**  
University of Silesia, Poland

Polynomial-coefficient generalizations of the Levi-Civita and Wilson  
functional equations  
(joint work with MACIEJ SABLIK)

THEOREM

If continuous functions  $f_1, \dots, f_M: \mathbb{R} \rightarrow \mathbb{C}$  satisfy the functional equation

$$\sum_{i=0}^M f_i(x + b_i y) P_i(x, y) = \sum_{j=1}^n u_j(x) v_j(y), \quad b_i \neq b_j \text{ for } i \neq j \quad (1)$$

with some polynomials  $P_i$  and some continuous functions  $u_j, v_j$ , then each  $f_i$  is a ratio of an exponential polynomial and a polynomial:

$$f_i(x) = \sum_{j=1}^n e^{\lambda_j x} r_j(x),$$

where  $r_j$  are rational functions.

We discuss also some generalizations of equation (1).

**Justyna Sikorska**  
University of Silesia, Poland

Various notions of orthogonality and the Cauchy functional equation

On the example of the famous Cauchy functional equation we show how various notions of orthogonality appear in the theory of functional equations. We give solutions of the Cauchy equation postulated for orthogonal vectors. Applications of this conditional equation both inside and outside mathematics constitutes a significant part of the lecture. Furthermore, we plan to discuss various aspects of stability problem. Last, but not least, some open problems concerning the topic will be presented.

# Slavko Simić

Mathematical Institute SANU, Serbia

## Stolarsky means in many variables

There is a huge amount of papers investigating properties of the so-called Stolarsky (or extended) two-parametric mean value, defined for positive values of  $x, y; x \neq y$ , as

$$E_{r,s}(x, y) := \left( \frac{r(x^s - y^s)}{s(x^r - y^r)} \right)^{1/(s-r)}, \quad rs(r-s) \neq 0.$$

Those means can be continuously extended on the domain  $\{(r, s; x, y) | r, s \in \mathbb{R}; x, y \in \mathbb{R}_+\}$  by the following

$$E_{r,s}(x, y) = \begin{cases} \left( \frac{r(x^s - y^s)}{s(x^r - y^r)} \right)^{1/(s-r)}, & rs(r-s) \neq 0; \\ \exp\left(-\frac{1}{s} + \frac{x^s \log x - y^s \log y}{x^s - y^s}\right), & r = s \neq 0; \\ \left( \frac{x^s - y^s}{s(\log x - \log y)} \right)^{1/s}, & s \neq 0, r = 0; \\ \sqrt{xy}, & r = s = 0; \\ x, & y = x > 0, \end{cases}$$

and in this form are introduced by Keneth Stolarsky in [1]. There are several papers attempting to define an extension of the class  $E$  to  $n$ ,  $n > 2$  variables [1].

In this talk we shall expose two possible explicit formulae of Stolarsky means in  $n$  variables which preserve its main properties and coincide for  $n = 2$ .

DEFINITION 1.

Let  $X_n = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$ . Then,

$$e_{r,s}(X_n) = e_{r,s}(x_1, x_2, \dots, x_n) := \left( \frac{r^2 x_1^{ns} + x_2^{ns} + \dots + x_n^{ns} - n(x_1 x_2 \dots x_n)^s}{s^2 x_1^{nr} + x_2^{nr} + \dots + x_n^{nr} - n(x_1 x_2 \dots x_n)^r} \right)^{\frac{1}{n(s-r)}}, \quad rs(s-r) \neq 0.$$

DEFINITION 2.

Let  $A_n = (a_1, a_2, \dots, a_n), X_n = (x_1, x_2, \dots, x_n), Y_n = (y_1, y_2, \dots, y_n); A_n, X_n, Y_n \in \mathbb{R}_+^n$ . Then,

$$E_{r,s}^n(A_n; X_n, Y_n) := \left( \frac{r^2 a_1(x_1^s - y_1^s)^2 + a_2(x_2^s - y_2^s)^2 + \dots + a_n(x_n^s - y_n^s)^2}{s^2 a_1(x_1^r - y_1^r)^2 + a_2(x_2^r - y_2^r)^2 + \dots + a_n(x_n^r - y_n^r)^2} \right)^{\frac{1}{2(s-r)}}.$$

Both extensions are symmetric and monotone increasing in both parameters  $r$  and  $s$  with  $e_{r,s}(x_1, x_2) = E_{r,s}^1(a_1; x_1, x_2) = E_{r,s}(x_1, x_2)$ .

### References

- [1] K.B. Stolarsky, *Generalizations of the logarithmic mean*, Math. Mag., 48(2)(1975), pp. 87-92.
- [2] J.K. Merikowski, *Extending means of two variables to several variables*, J. Ineq. Pure Appl. Math., 5(3) (2004), Article 65.
- [3] S. Simić, *On weighted Stolarsky means*, Sarajevo J. Math., Vol. 7(19) (2011).

## The short ruler on the general affine group

The restriction to the interval  $[0, 1]$  of a homomorphism  $h: (\mathbb{R}, +) \rightarrow (G, \circ)$  on a Lie group  $G$  is a geodesic. The problem is to construct long geodesics. We assume that we have a short ruler, which allows constructing geodesics with length  $L > 0$ . We can shorten a curve  $\alpha$  on  $G$  using the short ruler (reduced transformation).




Figure 1: The reduced transformation.

The reduced process  $(R_L^t \alpha)_{t \in \mathbb{N}}$  is the iteration of this transformation. In normed vector spaces, the reduced process converges to the straight line. On the general affine group  $\text{Aff}(1, \mathbb{R})$ —which is a Lie group—the reduced process converges to the geodesic linking the starting point of the curve  $\alpha$  with its end point.

### References

- [1] G.D. Birkhoff, *Dynamical Systems*. American Mathematical Society Colloquium Publications volume IX. Amsterdam: American Mathematical Society 1927. ISBN 978-0-8218-1009-5. doi:10.1090/co11/009.
- [2] J. Jost, *Riemannian Geometry and Geometric Analysis*. Universitext. Springer 1995. ISBN 0387571132. doi:10.1007/978-3-319-61860-9.
- [3] R. Liedl, N. Netzer, “Group Theoretic and Differential Geometric Methods for Solving the Translation Equation”. In *European Conference on Iteration Theory (ECIT 87)*, pp. 240–252. World Scientific Publishing 1989. ISBN 981-02-0041-2.
- [4] P. Stadler, “Curve Shortening by Short Rulers”. In *Journal of Difference Equations and Applications* **22** (1), pp. 22–36. London: Taylor & Francis 2016. ISSN 1563-5120. doi:10.1080/10236198.2015.1073724.

## Henrik Stetkær

Aarhus University, Denmark

### The Small Dimension Lemma revisited

Dilian Yang [Y] used the Small Dimension Lemma about irreducible, unitary representations of a compact group to solve d'Alembert's and Wilson's functional equations on such a group. We present a purely algebraic generalization of the Small Dimension Lemma. By help of it we find on any compact group  $G$  the solutions  $f, g \in C(G)$  of generalizations of d'Alembert's and Wilson's functional equations of the form

$$f(xy) + \mu(y)f(xy^*) = 2f(x)g(y), \quad x, y \in G,$$

where  $\mu \in C(G)$  is a given character of  $G$ , and  $x \mapsto x^*$  is a given, continuous involution of  $G$ .

#### References

[Y] D. Yang, *Functional equations and Fourier analysis*. *Canad. Math. Bull.* **56** (2013), 218–224.

## Tomasz Stypuła

Pedagogical University of Kraków, Poland

### Orthogonality preserving property on small sets

In this report we consider a problem when the orthogonality preserving property of a linear mapping on a small set, implies its orthogonality preserving property on the whole space. In order to construct such sets, we introduce a concept of independent bases. We present examples and results in finite-dimensional real inner product spaces.

## Mariusz Sudzik

University of Zielona Góra, Poland

### Iterative functional equations and attractive fixed points

Let  $I$  be a nontrivial interval and  $f, g : I \rightarrow I$  be given functions. We will consider the functional equation

$$\varphi(x) = \frac{\varphi(f(x)) + \varphi(g(x))}{2}$$

under the assumption that  $f$  and  $g$  have a globally attractive fixed point. Equations of higher order and their inhomogeneous versions will be analysed as well.

## László Székelyhidi

University of Debrecen, Hungary

### Functional equations on infinite joins

(joint work with ŻYWILLA FECHNER)

In this talk we present some results about basic function classes on infinite hypergroup joins. These results may serve as a starting point to study convolution type functional equations on infinite hypergroup joins using spectral synthesis.

## Imke Toborg

Martin-Luther-Universität Halle-Wittenberg, Germany

### On the functional equation $f(x)^{-1} = f^{-1}(x)$ on groups

In [1] David J. Schmitz introduced the notion of an inverse ambiguous function.

A bijective function from a group into itself is inverse ambiguous if and only if it is a solution of  $f(x)^{-1} = f^{-1}(x)$ . In this talk we give a precise description when a finite group admits an inverse ambiguous function or an inverse ambiguous automorphism.

#### References

- [1] D.J. Schmitz, *Inverse ambiguous functions on fields*, Aequat. Math. **91** (2017), 373–389.

## Anita Tomar

Government Degree College Raipur, India

### Fixed point and applications of Hardy-Rogers type contraction

In physical world a fixed point signifies a condition wherever a stable state or equilibrium is attained. Presence of fixed point plays a significant role in nonlinear analysis as numerous real-world problems in applied science, economics, chemistry, physics, computer science and engineering can be reformulated as a problem of finding fixed points of nonlinear maps. The aim of this talk is to discuss the existence of fixed point of almost  $\alpha$ -Hardy-Rogers- $\mathcal{F}$ -contraction in a partial metric space. Finally, some interesting examples and the solution of differential equation arising in critically damped harmonic oscillator is also discussed to demonstrate the usability of results.

#### References

- [1] M. Cosentino and P. Vetro, *Fixed point results for  $F$ -contractive mappings of Hardy-Rogers-type*, *Filomat*, **28**(4) (2014), 715–722.
- [2] S. G. Matthews, *Partial metric topology*, Research Report 212, Department of Computer Science, University of Warwick, 1992.
- [3] S. G. Matthews, *Partial metric topology*, Proceedings of the 8th Summer Conference on General Topology and Applications, Ann. New York Acad. Sci., **728** (1994), 183–197.
- [4] B. Samet, C. Vetro and P. Vetro, *Fixed point theorems for  $\alpha - \psi$ -contractive type mappings*, *Nonlinear Anal*, **75** (2012), 2154–2165.
- [5] D. Wardowski, *Fixed points of new type of contractive mappings in complete metric space*, *Fixed Point Theory Appl*, 2012.

## Andrzej Wiśnicki

Pedagogical University of Kraków, Poland

### Around the nonlinear Ryll-Nardzewski theorem

In this talk, we show that if  $S$  is a distal semigroup of nonexpansive mappings acting on a weak\* compact convex subset  $Q$  of a dual Banach space with the Radon-Nikodým property, then there is a common fixed point of  $S$  in  $Q$ . In particular, it gives a nonlinear counterpart of the Ryll-Nardzewski theorem. As a consequence, we obtain a nonlinear extension of the Bader-Gelander-Monod theorem concerning isometries in  $L$ -embedded Banach spaces.

#### References

- [1] U. Bader, T. Gelander, N. Monod, *A fixed point theorem for  $L_1$  spaces*, *Invent. Math.* **189** (2012), 143–148.
- [2] C. Ryll-Nardzewski, *Generalized random ergodic theorems and weakly almost periodic functions*, *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* **10** (1962), 271–275.
- [3] A. Wiśnicki, *Around the nonlinear Ryll-Nardzewski theorem*, arXiv:1903.12123.



# Alfred Witkowski

UTP University of Science and Technology, Poland

## Inequalities of Levin-Stečkin and Clausing

The Levin-Stečkin inequality comes from the Appendix to the Russian translation of “Inequalities” [3].

**THEOREM (LEVIN-STEČKIN’S INEQUALITY)**

If a function  $p: [0, 1] \rightarrow \mathbb{R}$  satisfies the conditions

- $p$  is non-decreasing in  $[0, 1/2]$ ,
- $p$  is symmetric, i.e.  $p(x) = p(1 - x)$ ,

then for every convex function  $\varphi$  the following inequality holds

$$\int_0^1 p(x)\varphi(x)dx \leq \int_0^1 p(x)dx \int_0^1 \varphi(x)dx.$$

In 1980 Clausing [2] proved the following theorem.

**THEOREM (CLAUSING’S INEQUALITY)**

Let  $p$  be a nonnegative functions on  $[0, 1]$  satisfying the following conditions:

- $p$  is non-decreasing on  $[0, 1/2]$ ,
- $p$  is symmetric.

Then for every concave, positive function  $\varphi$  the inequality

$$\int_0^1 p(x)\varphi(x)dx \leq \int_0^1 4 \min\{x, 1 - x\}p(x)dx \int_0^1 \varphi(x)dx$$

holds.

We provide new, elementary proofs of the above theorems.

## References

- [1] H. Brunn, *Nachtrag zu dem Aufsatz über Mittelwertssätze für bestimmte Integrale*, Münchener Berichte 1903, 205–212.
- [2] A. Clausing, *Disconjugacy and Integral Inequalities*, Trans. Amer. Math. Soc. **260** (1980), 293–307.
- [3] G.H. Hardy, J.E. Littlewood, G. Polya, *Inequalities*, Moscow, 1948 (in Russian)
- [4] V.I. Levin, S.B. Stečkin, *Inequalities*, Amer. Math. Soc. Transl. (2) **14** (1960), 1–29
- [5] P.R. Mercer, *A note on inequalities due to Clausing and Levin-Stečkin*, J. Math. Ineq. **11**(1) (2017), 163–166
- [6] A. Witkowski, *Inequalities of Levin-Stečkin, Clausing and Chebyshev revisited*, Elemente der Mathematik, to appear.

## Paweł Wójcik

Pedagogical University of Kraków, Poland

### Semi-smooth points in space $\mathcal{K}(H_1, H_2)$

The investigations of the smooth points in the operator spaces  $\mathcal{K}(H)$  were started in [1]. The aim of this report is to discuss a characterization of semi-smooth points in the compact operator space  $\mathcal{K}(H_1, H_2)$ , where  $H_1, H_2$  are Hilbert spaces.

#### References

- [1] J. R. Holub, *On the Metric Geometry of Ideals of Operators on Hilbert Space*, Math. Ann. **201** (1973), 157–163.
- [2] P.M. Miličić, *Sur le semi-produit scalaire dans quelques espaces vectoriel normés*, Mat. Vesnik, **8**(23) (1971), 181–185.

## Sebastian Wójcik

University of Rzeszów, Poland

### On convexity of the Swiss premium principle

*The Swiss premium principle* for a risk, represented by a non-negative bounded random variable  $X$  on a given probability space, is defined as a unique real number  $H_{u,c}(X)$  satisfying equation

$$u((c-1)H_{u,c}(X)) = E[u(cH_{u,c}(X) - X)] \quad (1)$$

where  $c \in [0, 1]$  and  $u : \mathbb{R} \rightarrow \mathbb{R}$  is a strictly increasing continuous function with  $u(0) = 0$  (cf. [1]). In the particular cases  $c = 0$  and  $c = 1$  the Swiss premium principle reduces to *the mean-value principle* and *the zero utility principle*, respectively.

In the talk, applying some results in [2, 3, 4], we present a characterization of convexity of the functional  $H_{u,c}$  defined implicitly by (1).

#### References

- [1] H. Bühlmann, B. Gagliardi, H. Gerber, E. Straub, *Some inequalities for stop-loss premiums*, ASTIN Bulletin **9** (1977), 75–83.
- [2] J. Chudziak, D. Głazowska, J. Jarczyk, W. Jarczyk, *On weighted quasi-arithmetic means which are convex*, Math. Inequal. Appl., in press.
- [3] Zs. Páles, *General inequalities for quasideviation means*, Aequationes Math. **36** (1988), 32–56.
- [4] Zs. Páles, P. Pasteczka, *On the best Hardy constant for quasi-arithmetic means and homogeneous deviation means*, Math. Inequal. Appl. **21** (2018), 585–599.

## Amr Zakaria

University of Debrecen, Hungary

### Equality problems related to Cauchy means

(joint work with ZSOLT PÁLES)

In this talk we establish a new characterization of the equality of two-variable Cauchy means (cf. [4]) to two-variable quasi-arithmetic means (cf. [1]) under natural, and therefore, the weakest possible regularity conditions. As an immediate application, we shed new light on the equality problem of two-variable Cauchy means which was solved by Losonczi [3, Theorem 5] under seven times differentiability assumptions. The approach is based on the results of the paper [2].

#### References

- [1] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge University Press, Cambridge, 1934, (first edition), 1952 (second edition).
- [2] T. Kiss and Zs. Páles, *On a functional equation related to two-variable weighted quasi-arithmetic means*, J. Difference Equ. Appl. **24** (2018), 107–126.
- [3] L. Losonczi, *Equality of two variable Cauchy mean values*, Aequationes Math. **65** (2003), 61–81.
- [4] E. Leach and M. Sholander, *Multivariable extended mean values*, J. Math. Anal. Appl. **104** (1984), 390–407.

## Thomas Zürcher

University of Silesia, Poland

### Around a Kazimierz Nikodem result - part II

(joint work with JANUSZ MORAWIEC)

This is joint work with Janusz Morawiec, and he will deliver part I. In the first part, equations of the form

$$\varphi(x) = \sum_{n=1}^N |f'_n(x)|\varphi(f_n(x)) + g(x)$$

will be considered. In this talk, we are changing the derivatives  $f'_n$  to some other functions  $g_n$ , looking for solutions  $\varphi \in L^1([0, 1])$  of

$$\varphi(x) = \sum_{n=1}^N |g_n(x)|\varphi(f_n(x)) + g(x).$$

This is not only a cosmetic change. We need new methods to tackle this kind of equations.

**Marcin J. Zygmunt**  
University of Silesia, Poland

Additive functions on "ax+b" group

The aim of the talk is to solve Pexider's equation  $f(xy) = g(x)h(y)$  for functions  $f, g, h : G \rightarrow G$  acting on a noncommutative group  $G$ . The equation will be completely solved in the case of affine group "ax+b".