



LIST OF TALKS

1. ANA MARIA ACU 🌐, *Applications of Ohlin lemma and Levin-Stečkin inequalities*
2. MIROSLAW ADAMEK, *Some remarks about convex sequences*
3. JAVID ALI 🌐, *Fixed Point Iterations: Convergence, stability and data dependence results*
4. LJILJANA ARAMBAŠIĆ 🌐, *Strong Birkhoff–James orthogonality in commutative C^* -algebras*
5. KAROL BARON, *Strong law of large numbers for iterates of some random-valued functions*
6. CHAIMAA BENZAROUALA 🌐, *A fixed point approach to the Ulam-Hyers-Rassias stability of a generalized linear functional equation*
7. ZOLTÁN BOROS 🌐, *Conditionally linked monomial functions*
8. PÁL BURAI, *Random means generated by random variables: expectation and limit theorems*
9. GOPALAKRISHNA CHAITANYA 🌐, *Square roots of functions*
10. JACEK CHMIELIŃSKI, *On continuity of seminorms*
11. JACEK CHUDZIAK, *On existence and properties of the principle of equivalent utility*
12. BEATA DERĘGOWSKA, *Quasiarithmetic-type invariant means on probability space*
13. HOJJAT FARZADFARD 🌐, *On the Real Poincaré Functional Equation*
14. ŻYWILLA FECHNER 🌐, *Moment functions and exponential monomials on commutative hypergroups*
15. GIAN LUIGI FORTI, *Alternative Cauchy equation in three unknown functions*
16. ROMAN GER, *On alienation of two functional equations of quadratic type (again)*
17. ATTILA GILÁNYI, **Computer assisted studies of functional equations and inequalities (L)**
18. DOROTA GLAZOWSKA, *Generalized classical weighted means and some related problems*
19. KAROL GRYSZKA 🌐, *On the divergence of continuous functions*
20. ESZTER GSELMANN 🌐, *Polynomial identities satisfied by generalized polynomials*
21. GRZEGORZ GUZIK, *On geometric rate of convergence for generalized random iterations*
22. LÁSZLÓ HORVÁTH 🌐, *Refinements of the integral Jensen's inequality generated by finite or infinite permutations*
23. ELIZA JABŁOŃSKA, *Remarks on classes $\mathcal{A}, \mathcal{B}, \mathcal{C}$*
24. JUSTYNA JARCZYK, *On generalized Archimedes–Borchardt algorithm*
25. WITOLD JARCZYK, *Form of semiflows of pairs of weighted quasi-arithmetic means*
26. SÁNDOR JENEI, *Algebraic methods in functional equations – beyond the regularity conditions*

27. SOON-MO JUNG 🌱, **Approximate isometries on bounded sets (L)**
28. GERGELY KISS, *Charaterization of quasi-arithmetic means without regularity*
29. GRZEGORZ KLESZCZ, *On a functional–difference inclusion induced by some iterated function system*
30. TOMASZ KOCHANEK, *Approximately order zero maps between C^* -algebras*
31. ZBIGNIEW LEŚNIAK 🌱, *On a construction of iterative roots of a Brouwer homeomorphism*
32. RADOSŁAW ŁUKASIK, *Vector-valued invariant means and projections from the bidual space*
33. ARPITA MAL 🌱, *Extreme contraction from the perspective of multi-smoothness*
34. DANIELA MARIAN 🌱, *On Hyers-Ulam-Rassias stability of certain partial differential equations and integral equations*
35. JOANNA MARKOWICZ, *Uniform convexity of direct sums and interpolation spaces*
36. JANUSZ MATKOWSKI 🌱, *Remarks on fixed point theorem*
37. RAYENE MENZER 🌱, *Alternative equations for quadratic functions*
38. FLAVIA-CORINA MITROI-SYMEONIDIS, *Redistributing algorithms and Shannon’s Entropy*
39. MĀDĀLINA MOGA, *On some properties of Meir-Keeler operators*
40. LAJOS MOLNÁR 🌱, *On the order determining property of the norm of a Kubo-Ando mean in operator algebras*
41. VEERAPAZHAM MURUGAN 🌱, *Iterative root problem for continuous functions of non-monotonicity height 1*
42. HEMANT KUMAR NASHINE 🌱, *Fixed point results on relational metric spaces with applications to Hyers-Ulam stability and nonlinear matrix equations*
43. KAZIMIERZ NIKODEM, *Remarks on set-valued means*
44. CHISOM PRINCE OKEKE 🌱, *On some new functional equations characterizing polynomial functions*
45. ANDRZEJ OLBRYŚ, *On characterization of affine differences and related forms*
46. ADAM OSTASZEWSKI 🌱, *Tilted solutions of the Gotȃb-Schinzel functional equation in Banach algebras*
47. ZDZISŁAW OTACHEL, *New advances on some inequalities in inner product spaces*
48. ZSOLT PÁLES 🌱, *Limit theorems for deviation means of independent and identically distributed random variables*
49. PAWEŁ PASTECZKA, *On the Riemannian approach to integral means*
50. KALLOL PAUL 🌱, *k -smoothness of operators on infinite-dimensional spaces*
51. ADRIAN PETRUŞEL, **Existence and Stability Results in Fixed Point Theory (L)**
52. DORIAN POPA 🌱, *Best constant in Ulam Stability*
53. TERESA RAJBA, *On the Raşa inequality for higher-order convex functions*
54. IOAN RAŞA 🌱, *A functional equation related to Appell polynomials and Heun functions*

55. LENKA RUCKÁ, *Some results on distributionally chaotic points*
56. MACIEJ SABLIK, *On some new functional equations characterizing polynomial functions*
57. DEBMALYA SAIN 🌐, *Best approximations from the perspective of orthogonality*
58. ADOLF SCHLEIERMACHER, *On even orthogonally additive mappings*
59. JUAN MATÍAS SEPULCRE 🌐, *A class of functional equations associated with almost periodic functions*
60. YONG-GUO SHI 🌐, *Periodic points of asymmetric Bernoulli shifts*
61. JUSTYNA SIKORSKA, *On a generalized n -linear equation*
62. SLAVKO SIMIĆ 🌐, *Global bounds for the generalized Jensen functional with applications*
63. LÁSZLÓ SZÉKELYHIDI, *Functional equations, moment functions and Bell polynomials*
64. PATRICIA SZOKOL, *Hermite - Hadamard type inequality for certain Schur convex functions*
65. TOMASZ SZOSTOK, *On some class of means*
66. LAN NHI TO, *Computer assisted investigation of the alienness of linear functional equations*
67. RADU TRUȘCĂ, *Some local fixed point results and applications for generalized contractions*
68. FILIP TUROBOŚ 🌐, *On preserving the relaxed polygonal inequality*
69. SZYMON WĄSOWICZ 🌐, *New adaptive method of the approximate integration*
70. BETTINA WILKENS, *Algebraic methods in spectral synthesis on discrete abelian groups*
71. PAWEŁ WÓJCIK, *Generalized orthogonality equation*
72. SEBASTIAN WÓJCIK, *Quasi-arithmetic type means generated by the generalized Choquet integral*
73. AMR ZAKARIA 🌐, *An invariance problem with six unknown functions*
74. MAREK CEZARY ZDUN, *On a linearization of Koopmans recursion and its application in economics*
75. WEINIAN ZHANG 🌐, **Iteration and Equations (L)**
76. THOMAS ZÜRCHER, *New solutions to the Matkowski–Wesołowski problem*

L – Lecture 45 minutes, 🌐 – online participants.



ABSTRACTS OF TALKS

Ana Maria Acu 

Lucian Blaga University of Sibiu, Romania

Applications of Ohlin lemma and Levin-Stečkin inequalities (joint work with Ioan Raşa)

Let $n \geq 1$ and $a_{n,j} := 4^{-n} \binom{2j}{j} \binom{2n-2j}{n-j}$, $j = 0, 1, \dots, n$. The probability distribution $(a_{n,j})_{j=0,\dots,n}$ is involved in the study of some entropies: see [1], where two proofs of the inequality

$$\frac{1}{n+1} \sum_{j=0}^n f(j) \leq \sum_{j=0}^n a_{n,j} f(j), \quad f: \mathbb{R} \rightarrow \mathbb{R} \text{ convex}, \quad (1)$$

are given. One proof is based on the Ohlin Lemma. In this talk we discuss the connection of (1) with the Levin-Stečkin inequalities. New inequalities are obtained for strongly convex functions, combining (1) with results from [2]. Finally, we obtain estimates for the information potential associated with the distribution $(a_{n,j})_{j=0,\dots,n}$.

References

- [1] Acu, Ana Maria, Gulen Başcanbaz-Tunca, and Ioan Raşa. "Bounds for indices of coincidence and entropies." *Math. Inequal. Appl.* 24, no. 2 (2021): 307-321.
- [2] Nikodem, Kazimierz, and Teresa Rajba. "Ohlin and Levin-Stečkin-type results for strongly convex functions." *Ann. Math. Sil.* 34, no. 1 (2020): 123-132.

Mirosław Adamek
University of Bielsko-Biala, Poland

Some remarks about convex sequences

Let Z be a set of consecutive integers with $|Z| \geq 3$. A sequence of real numbers $(a_n)_{n \in Z}$ is called convex (see [2]), if it satisfies the following inequality

$$a_n \leq \frac{a_{n-1} + a_{n+1}}{2}$$

for all $n - 1, n, n + 1 \in Z$.

In this talk we will recall some known facts about convex sequences and give a discrete version of the result presented in [1], i.e. we will give a necessary and sufficient condition for the separation of two sequences with a convex sequence. As a consequence of a separation theorem we will obtain Hyers–Ulam stability type result for convex sequences. We will also give, as a consequence of the presented results for convex sequences, results concerning generalized convex sequences in the sense of [3].

References

- [1] Baron, Karol, Janusz Matkowski, and Kazimierz Nikodem. "A sandwich with convexity." *Math. Pannon.* 5, no. 1 (1994): 139-144.
- [2] Mitrinović, Dragoslav S., and Petar M. Vasić. *Analytic Inequalities*. New York: Springer-Verlag, 1970.
- [3] Tabor, Jacek, Józef Tabor, and Marek Źołdak. "Strongly convex sequences." In: *Inequalities and applications 2010*, 183-188. Vol. 161 of *International Series of Numerical Mathematics*. Basel: Birkhäuser, 2012.

Javid Ali 

Aligarh Muslim University, India

Fixed Point Iterations: Convergence, stability and data dependence results

In this talk we discuss a newly introduced two step fixed point iterative algorithm. We prove, under certain conditions, a strong convergence result for weak contractions $T: C \rightarrow C$, where C is a nonempty, closed and convex subset of a Banach space. We also prove stability and data dependency of a proposed iterative algorithm. Furthermore, we utilize our main result to approximate the solution of a nonlinear functional Volterra integral equation.

References

- [1] Ali, Javid, Mohd Jubair, and Faeem Ali. "Stability and convergence of F iterative scheme with an application to the fractional differential equation". *Engg. with Comp.* (2020): Online published.
- [2] Berinde, Vasile. "Summable almost stability of fixed point iteration procedures." *Carpathian J. Math.* 19, no. 2 (2003): 81-88.
- [3] Berinde, Vasile. *Iterative approximation of fixed points*. Vol. 1912 of *Lecture Notes in Mathematics*. Berlin Heidelberg: Springer-Verlag, 2007.
- [4] Harder, Alberta Marie. Fixed point theory and stability results for fixed point iteration procedures. Vol. 634 of *Doctoral Dissertations*. University of Missouri–Rolla, 1987.
- [5] Rus, Ioan A., and Sorin Mureşan. "Data dependence of the fixed points set of weakly Picard operators." *Studia Univ. Babeş-Bolyai Math.* 43, no. 1 (1998): 79-83.

Ljiljana Arambašić 

University of Zagreb, Croatia

Strong Birkhoff–James orthogonality in commutative C^* -algebras (joint work with Alexander Guterman, Bojan Kuzma, Rajna Rajić, and Svetlana Zhilina)

General normed spaces are not equipped with an inner product. Nonetheless, there do exist several nonequivalent extensions of orthogonality from inner product spaces to general normed ones. One of the most well-known is the Birkhoff–James orthogonality: if X is a normed space and $x, y \in X$, then x is Birkhoff–James orthogonal to y if $\|x + \lambda y\| \geq \|x\|$ holds for all scalars λ . In a C^* -algebra A we can also discuss the case when $x \in A$ is Birkhoff–James orthogonal to all the elements of the form ya , $a \in A$. In this case we say that x is strongly Birkhoff–James orthogonal to y . We discuss this kind of orthogonality in commutative C^* -algebras.

Karol Baron

University of Silesia, Katowice, Poland

Strong law of large numbers for iterates of some random-valued functions (joint work with Rafał Kapica)

Fix a probability space (Ω, \mathcal{A}, P) and a metric space X . Let \mathcal{B} denote the σ -algebra of all Borel subsets of X . We say that $f: X \times \Omega \rightarrow X$ is a *random-valued function* (an *rv-function* for short) if it is measurable with respect to the product σ -algebra $\mathcal{B} \otimes \mathcal{A}$. The iterates of such an rv-function are given by

$$f^0(x, \omega_1, \omega_2, \dots) = x, \quad f^n(x, \omega_1, \omega_2, \dots) = f(f^{n-1}(x, \omega_1, \omega_2, \dots), \omega_n)$$

for $n \in \mathbb{N}$, $x \in X$ and $(\omega_1, \omega_2, \dots)$ from Ω^∞ defined as $\Omega^\mathbb{N}$. Note that $f^n: X \times \Omega^\infty \rightarrow X$ is for every $n \in \mathbb{N}$ an rv-function on the product probability space $(\Omega^\infty, \mathcal{A}^\infty, P^\infty)$. See [4], Sec. 1.4, and [2].

Many results on the convergence of such iteration sequences can be found in the literature. They concern different type of convergence and are given under suitable assumptions concerning both the metric space and the rv-function. We accept the following hypothesis:

(H) (X, ρ) is a complete and separable metric space and $f: X \times \Omega \rightarrow X$ is an rv-function such that

$$\int_{\Omega} \rho(f(x, \omega), f(z, \omega)) P(d\omega) \leq \lambda \rho(x, z) \quad \text{for } x, z \in X$$

with a $\lambda \in (0, 1)$, and

$$\int_{\Omega} \rho(f(x, \omega), x) P(d\omega) < \infty \quad \text{for } x \in X.$$

Then (see [1] and [3]) there exists a probability Borel measure π^f on X such that for every $x \in X$ the sequence of distributions of $f^n(x, \cdot)$, $n \in \mathbb{N}$, converges weakly to π^f and

$$\int_X \rho(x, z) \pi^f(dz) < \infty.$$

We are looking for conditions that imposed on the rv-function $f: X \times \Omega \rightarrow X$ and a function $\psi: X \rightarrow \mathbb{R}$ provide that for every $x \in X$ we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \psi \circ f^k(x, \cdot) = \int_X \psi d\pi^f \quad \text{a.e. for } P^\infty.$$

References

- [1] Baron, Karol. "On the convergence in law of iterates of random-valued functions." *Aust. J. Math. Anal. Appl.* 6, no. 1 (2009): Art. 3.
- [2] Kapica, Rafał. "Sequences of iterates of random-valued vector functions and solutions of related equations." *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 213 (2004): 113-118 (2005).
- [3] Kapica, Rafał. "The geometric rate of convergence of random iteration in the Hutchinson distance." *Aequationes Math.* 93, no. 1 (2019): 149-160.
- [4] Kuczma, Marek, Bogdan Choczewski, and Roman Ger. *Iterative Functional Equations*. Vol. 32 of *Encyclopedia of Mathematics and its Applications*. Cambridge: Cambridge Univ. Press, 1990.

Chaimaa Benzarouala 
Mohammed V University in Rabat, Morocco

A fixed point approach to the Ulam-Hyers-Rassias stability of a generalized
linear functional equation
(joint work with Lahbib Oubbi)

During this talk, we introduce the general functional equation

$$\sum_{i=1}^m A_i(f(\varphi_i(\bar{x}))) + b = 0, \quad \bar{x} := (x_1, \dots, x_n) \in X^n,$$

and study its Ulam-Hyers-Rassias stability and hyperstability, using a fixed point approach, where m and n are positive integers, f is a mapping from a vector space X into a Banach space $(Y, \|\cdot\|)$, and for every $i \in \{1, \dots, m\}$, φ_i is a linear mapping from X^n into X , A_i is a continuous endomorphism of Y and $b \in Y$. Our result covers most of the former ones in the literature concerning the stability and hyperstability of linear functional equations, as well as new situations. This talk is based on the article [3].

References

- [1] Bahyrycz, Anna, and Jolanta Olko. "On stability of the general linear equation." *Aequationes Math.* 89, no. 6 (2015): 1461-1474.
- [2] Bahyrycz, Anna, and Jolanta Olko. "Hyperstability of general linear functional equation." *Aequationes Math.* 90, no. 3 (2016): 527-540.
- [3] Benzarouala, Chaimaa, and Lahbib Oubbi. "Ulam-stability of a generalized linear functional equation, a fixed point approach." *Aequationes Math.* 94, no. 5 (2020): 989-1000.
- [4] Brzdęk, Janusz, Jacek Chudziak, and Zsolt Páles. "A fixed point approach to stability of functional equations." *Nonlinear Anal.* 74, no. 17 (2011): 6728-6732.
- [5] Zhang, Dong. "On hyperstability of generalised linear functional equations in several variables." *Bull. Aust. Math. Soc.* 92, no. 2 (2015): 259-267.
- [6] Oubbi, Lahbib. "Ulam-Hyers-Rassias stability problem for several kinds of mappings." *Afr. Mat.* 24, no. 4 (2013): 525-542.

Zoltán Boros 

University of Debrecen, Hungary

Conditionally linked monomial functions
(joint work with Edit Garda-Mátyás)

Among others, we establish the following generalized versions of our recent statements [2].

Theorem 1. *Suppose that $f, g: \mathbb{R} \rightarrow \mathbb{R}$ are generalized monomials of degree $n \in \mathbb{N}$ that satisfy the additional equation $y^n f(x) = x^n g(y)$ under the condition $y = a_m x^m + a_{m-1} x^{m-1} + \cdots + a_1 x + a_0$, where $m \in \mathbb{N}$, $a_i \in \mathbb{R}$ ($i = 0, \dots, m$) are fixed such that $a_m \neq 0$ and $a_0 \neq 0$. Then $f(x) = g(x) = x^n f(1)$ for all $x \in \mathbb{R}$.*

Theorem 2. *Let $n \in \{1, 2, 3\}$. If $m \in \mathbb{Z}$, $|m| \geq 2$ and $f: \mathbb{R} \rightarrow \mathbb{R}$ is a generalized monomial of degree n that satisfies the additional equation $y^n f(x) = x^n f(y)$ under the condition $y = x^m$, then $f(x) = x^n f(1)$ for all $x \in \mathbb{R}$.*

Our former results [1] for the case $n = 2$ indicate that we cannot immediately extend the latter statement for two generalized monomials in the conditional equation.

Further similar problems, statements and counterexamples are presented as well.

References

- [1] Boros, Zoltán, and Edit Garda-Mátyás. "Conditional equations for quadratic functions." *Acta Math. Hungar.* 154, no. 2 (2018): 389-401.
- [2] Boros, Zoltán, and Edit Garda-Mátyás. "Conditional equations for monomial functions." *Publ. Math. Debrecen* (accepted).

Pál Burai

Budapest University of Technology and Economics, Hungary


Random means generated by random variables: expectation and limit theorems

(joint work with Mátyás Barczy)

We introduce the notion of a random mean generated by a random variable and give a construction of its expected value. We derive some sufficient conditions under which strong law of large numbers and some limit theorems hold for random means generated by the elements of a sequence of independent and identically distributed random variables.

References

- [1] Baron, Karol. "Weak law of large numbers for iterates of random-valued functions." *Aequationes Math.* 93, no. 2 (2019): 415-423.
- [2] Baron, Karol, and Witold Jarczyk. "Random-valued functions and iterative functional equations." *Aequationes Math.* 67, no. 1-2 (2004): 140-153.
- [3] Beliakov, Gleb, Humberto Bustince Sola, and Tomasa Calvo Sánchez. *A practical guide to averaging functions*. vol. 329 of *Studies in Fuzziness and Soft Computing*. Cham: Springer, 2016.
- [4] Borsato, Luísa, Eduardo Horta, and Rafael Rigão Souza. "A characterization of the strong law of large numbers for Bernoulli sequences." *arXiv* (2020): arxiv.org/pdf/2008.00318.pdf.
- [5] Cohn, Donald L. *Measure theory*. Boston, Mass.: Birkhäuser, 1980.
- [6] Hytönen, Tuomas, Jan van Neerven, Mark Veraar, and Lutz Weis. *Analysis in Banach spaces. Vol. I. Martingales and Littlewood-Paley theory*, vol. 63 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Cham: Springer, 2016.
- [7] Jarczyk, Justyna, and Witold Jarczyk. "Invariance of means." *Aequationes Math.* 92, no. 5 (2018): 801-872.
- [8] Jarczyk, Justyna, and Witold Jarczyk. "Gaussian iterative algorithm and integrated automorphism equation for random means." *Discrete Contin. Dyn. Syst.* 40, no. 12 (2020): 6837-6844.
- [9] Kallenberg, Olav. *Random measures, theory and applications*, vol. 77 of *Probability Theory and Stochastic Modelling*. Cham: Springer, 2017.
- [10] Kuo, Hui Hsiung. *Gaussian measures in Banach spaces*. vol. 463 of *Lecture Notes in Mathematics*. Berlin-New York: Springer-Verlag, 1975.
- [11] Páles, Zsolt, and Amr Zakaria. "On the local and global comparison of generalized Bajraktarević means." *J. Math. Anal. Appl.* 455, no. 1 (2017): 792-815.
- [12] Panaretos, Victor M., and Yoav Zemel. *An invitation to statistics in Wasserstein space*. Cham: Springer, 2020.
- [13] Villani, Cédric. *Topics in optimal transportation*. Providence, RI: American Mathematical Society (AMS), 2003.
- [14] Wong, Willie. "Continuity of the infimum." *Bubbles Bad; Ripples Good, Blog*, 2011.

Gopalakrishna Chaitanya 

Indian Statistical Institute, Bangalore, India

Square roots of functions

An *iterative square root* of a function f is a function g such that $g(g(\cdot)) = f(\cdot)$. We discuss a new result on the nonexistence of iterative square roots of self-maps on arbitrary sets and use it to show that continuous self-maps without iterative square roots are dense in the space of all continuous self-maps on the unit cube in \mathbb{R}^m . This talk is based on a joint work with B. V. Rajarama Bhat. (see <https://arxiv.org/abs/2105.02171>).

Jacek Chmieliński

Pedagogical University of Krakow, Poland

On continuity of seminorms

(joint work with Moshe Goldberg)

Let X be an infinite-dimensional real or complex vector space. We study the continuity and discontinuity properties of seminorms with respect to norm-topologies on X . In particular, the following problems are considered:

- the existence of a norm on X , with respect to which a given seminorm is continuous/discontinuous;
- the existence of a norm on X , with respect to which all seminorms are continuous/discontinuous;
- the equivalence of norms, such that a given seminorm is continuous/discontinuous with respect to all of them.

References

- [1] Goldberg, Moshe. "Continuity of seminorms on finite-dimensional vector spaces." *Linear Algebra Appl.* 515 (2017): 175-179.
- [2] Chmieliński, Jacek, and Moshe Goldberg. "Continuity and discontinuity of seminorms on infinite-dimensional vector spaces." *Linear Algebra Appl.* 578 (2019): 153-158.
- [3] Chmieliński, Jacek, and Moshe Goldberg. "Continuity and discontinuity of seminorms on infinite-dimensional vector spaces. II." *Linear Algebra Appl.* 594 (2020): 249-261.

Jacek Chudziak

University of Rzeszów, Poland

On existence and properties of the principle of equivalent utility

The principle of equivalent utility is one of the methods of insurance contract pricing. If $w \in \mathbb{R}$ is an initial wealth of the insurance company and $u : \mathbb{R} \rightarrow \mathbb{R}$ is its continuous and strictly increasing utility function, then the premium of equivalent utility for a risk X , represented by a non-negative bounded random variable on a given probability space, is defined as a solution $H_{(w,u)}(X)$ of the equation

$$E[u(w + H_{(w,u)}(X) - X)] = u(w). \quad (1)$$

Several results concerning the principle defined by (1) can be found e.g. in Bowers et al. [1], Bühlmann [2], Gerber [3] and Rolski et al. [6].

Recently, the principle of equivalent utility has been extended onto various models of decision making under risk. Heilpern [4] proposed and investigated the principle under rank-dependent utility model. Kałuska and Krzeszowiec [5] introduced the principle of equivalent utility based on Cumulative Prospect Theory. In this setting the principle is defined through the equation

$$E_{gh} \left[u(w + H_{(w,u,g,h)}(X) - X) \right] = u(w), \quad (2)$$

where E_{gh} is the the generalized Choquet integral with respect to to the probability weighting functions g (for gains) and h (for losses) – cf. [5].

In the talk we establish a necessary and sufficient condition for the existence and uniqueness of the principle defined by (2). Furthermore we provide the characterizations of some important properties of the principle.

References

- [1] Bowers, Newton Lynn, Hans-Ulrich Gerber, James Charles Hickman et al. *Actuarial Mathematics*. Itasca, Illinois: The Society of Actuaries, 1986.
- [2] Bühlmann, Hans. *Mathematical Models in Risk Theory*. Berlin: Springer, 1970.
- [3] Gerber, Hans-Ulrich. *An Introduction to Mathematical Risk Theory*. S. S. Huebner Foundation, Homewood Illinois: R.D. Irwin Inc., 1979
- [4] Heilpern, Stanisław. "A rank-dependent generalization of zero utility principle." *Insurance Math. Econom.* 33, no. 1 (2003): 67-73.
- [5] Kałuska, Marek, and Michał Krzeszowiec. "Pricing insurance contracts under Cumulative Prospect Theory." *Insurance Math. Econom.* 50, no. 1 (2012): 159-166.
- [6] Rolski, Tomasz, Hanspeter Schmidli, Volker Schmidt, et. al. *Stochastic Processes for Insurance and Finance*. New York: John Wiley & Sons, 1999.

Beata Deręgowska
Pedagogical University of Krakow, Poland

Quasiarithmetic-type invariant means on probability space
(joint work with Paweł Pasteczka)

For a family $(\mathcal{A}_x)_{x \in (0,1)}$ of integral quasi-arithmetic means satisfying certain measurability-type assumptions we search for an integral mean K such that $K\left((\mathcal{A}_x(\mathbb{P}))_{x \in (0,1)}\right) = K(\mathbb{P})$ for every compactly supported probability Borel measure \mathbb{P} . Also some results concerning the uniqueness of invariant means will be given.

References

- [1] Deręgowska, Beata, and Paweł Pasteczka. "Quasi-arithmetic-type invariant means on probability space." *Aequationes Math.* 95, no. 4 (2021): 639-651.

On the Real Poincaré Functional Equation

Let I be a real interval, $f : I \rightarrow I$ and $\alpha \in \mathbb{R}$. A real Poincaré functional equation is of the form

$$\psi(\alpha x) = f(\psi(x)), \quad (x \in I),$$

where a function $\psi : \mathbb{R} \rightarrow I$ is unknown. We deal with the case where I is a compact interval and a continuous periodic solution ψ is desired. This talk is based on the papers [1] and [2]. The main results are as follows:

We call a function $\psi : \mathbb{R} \rightarrow \mathbb{R}$ a **cosine-like function** provided that ψ is continuous, nonconstant and periodic, moreover, if λ is the principal period of ψ , then

1. ψ is even, in particular $\psi(x) = \psi(\lambda - x)$ for all $0 \leq x \leq \lambda$.
2. If x and y are two distinct points in the interval $[0, \lambda]$ and $\psi(x) = \psi(y)$, then $y = \lambda - x$.
3. ψ attains one of its absolute extrema at 0 and the other at $\lambda/2$.
4. ψ is strictly increasing on one of the intervals $[0, \lambda/2]$ and $[\lambda/2, \lambda]$ and strictly decreasing on the other.

Let I be a compact interval of positive length, $f : I \rightarrow I$ and $k \in \mathbb{N}$. We say that f is a **fluctuating function of rank k** provided that there exists a partition $a_0 < a_1 < \dots < a_k$ of I such that for each $0 \leq j < k$ the function f is strictly monotone on the interval $[a_j, a_{j+1}]$ and $f([a_j, a_{j+1}]) = I$. The **node vector** of f is denoted by $\text{Node}(f) = (a_0, \dots, a_k)$. Also

$$\text{Node}_\infty(f) := \bigcup_{n=1}^{\infty} \text{Node}(f^n).$$

Theorem 1 (The Existence Theorem). *Let $I = [a, b]$ be a compact real interval and $f : I \rightarrow I$ be a fluctuating function of rank $k > 1$. The following statements are equivalent:*

1. *For every $\lambda > 0$ the equation $\psi(kx) = f(\psi(x))$ has a solution ψ which is a cosine-like function with the principal period λ .*
2. *$\text{Node}_\infty(f)$ is dense in I .*

Theorem 2 (The Uniqueness Theorem). *Let $\psi : \mathbb{R} \rightarrow I$ and $\phi : \mathbb{R} \rightarrow I$ be two solutions of the Poincaré functional equation $\psi(kx) = f(\psi(x))$ with ψ cosine-like and ϕ non-constant, continuous and periodic, where $k > 1$ is an integer. Then there exist a positive real number ω and an integer m such that $\phi(x) = \psi(\omega x + m\lambda/(k-1))$, ($x \in \mathbb{R}$), where λ is the principal period of ψ .*

References

- [1] Farzadfard, Hojjat. "Periodic solutions of the Poincaré functional equation: existence." *J. Math. Anal. Appl.* 498, no. 1 (2021): Art. no. 124933.
- [2] Farzadfard, Hojjat. "Periodic solutions of the Poincaré functional equation: uniqueness." *Proc. Amer. Math. Soc.* (2021). <https://doi.org/10.1090/proc/15682>.

Żywilla Fechner 

Lodz University of Technology, Poland

Moment functions and exponential monomials on commutative hypergroups (joint work with Eszter Gselmann and László Székelyhidi)

In this talk we are going to discuss a connection between moment functions and exponential monomials on commutative hypergroups. At the beginning we are going to give a brief motivation of the problem. In particular, we formalize the notions of an exponential monomial and moment functions. We want to find conditions under which exponential monomials can be expressed in terms of generalized moment functions. We present one of possible conditions and discuss some possible ways for further researchs.

References

- [1] Fechner, Żywilla, Eszter Gselmann, and László Székelyhidi. "Moment functions and exponential monomials on commutative hypergroups." *Aequat. Math.*, to appear.

Gian Luigi Forti

Università degli Studi di Milano, Italy

Alternative Cauchy equation in three unknown functions

In this paper we deal with the product of two or three Cauchy differences equaled to zero. We show that in the case of two Cauchy differences, the condition of absolute continuity and differentiability of the two functions involved implies that one of them must be linear, i.e. we have a trivial solution. In the case of the product of three Cauchy differences the situation changes drastically: there exists non trivial \mathcal{C}^∞ solutions, while in the case of real analytic functions we obtain that at least one of the functions involved must be linear. Some open problems are presented.

Roman Ger

University of Silesia, Katowice, Poland

On alienation of two functional equations of quadratic type (again)

We deal with an alienation problem for an Euler-Lagrange type functional equation

$$f(\alpha x + \beta y) + f(\alpha x - \beta y) = 2\alpha^2 f(x) + 2\beta^2 f(y)$$

assumed for fixed nonzero real numbers α, β , $1 \neq \alpha^2 \neq \beta^2$, and the classic quadratic functional equation

$$g(x + y) + g(x - y) = 2g(x) + 2g(y).$$

We were inspired by papers of Chang Il Kim, Giljun Han & Seong-A. Shim [3] and M.Eshagi Gordji & H. Khodaei [2], where the special case $g = \gamma f$ was examined, as well as by a suggestion of Bessenyei Mihály [1].

References

- [1] Bessenyei, Mihály. "Oral question during the discussion at the 20th Debrecen–Katowice Winter Seminar on Functional Equations and Inequalities, Hajdúszoboszló, Hungary, January 29 – February 1, 2020".
- [2] Eshaghi Gordji, Madjid, and Hamid Khodaei. "On the generalized Hyers-Ulam-Rassias stability of quadratic functional equations." *Abstr. Appl. Anal.* 2009: Article ID 923476.
- [3] Kim, Chang Il, Gil-Jun Han, and Seong-A. Shim. "Hyers-Ulam stability for a class of quadratic functional equations via a typical form." *Abstr. Appl. Anal.* 2013: Article ID 283173.

Attila Gilányi

University of Debrecen, Hungary

Computer assisted studies of functional equations and inequalities

During the last 30 years, computer assisted methods were applied in investigations of functional and inequalities by several researchers. Among others, solutions of general two variable linear functional equations were determined and further studies of linear equations were performed by computer, regularity properties of certain functional equations were tested, furthermore, various classes of means were studied with the help of such methods. In this talk, we present a systematic survey of these results.

Dorota Głazowska

University of Zielona Góra, Poland

Generalized classical weighted means and some related problems (joint work with Janusz Matkowski)

Under some simple conditions on real function f defined on an interval I , the two-variable functions given by the following formulas

$$A_f(x, y) := f(x) + y - f(y), \quad G_f(x, y) := \frac{f(x)}{f(y)} y \quad \text{and} \quad H_f(x, y) := \frac{xy}{f(x) + y - f(y)},$$

for all $x, y \in I$, generalize, respectively, the classical weighted arithmetic, geometric and harmonic means. In fact these means are symmetric, if and only if they coincide with $\mathcal{A}, \mathcal{G}, \mathcal{H}$, respectively. The invariance identity

$$\mathcal{G} \circ (A_f, H_f) = \mathcal{G},$$

extending the Pythagorean harmony proportion and confirming the adequacy of the generalized means, allows to conclude the suitable complementariness of A_f and H_f with respect to \mathcal{G} , and determine the convergence of sequence of the iterates of the mean-type mapping (A_f, H_f) to $(\mathcal{G}, \mathcal{G})$. Moreover, for each of the classical symmetric means $\mathcal{A}, \mathcal{H}, \mathcal{G}$ and for the above generalized means A_f, G_f, H_f we prove the existence and uniqueness of the respective complementary mean, we give its explicit formula, as well as the limit of the sequence of iterates of the relevant mean-type mappings. We also give solutions of some special cases of the invariance equation involving the above generalized classical weighted means.

Karol Gryszka

Pedagogical University of Krakow, Poland

On the divergence of continuous functions (joint work with Paweł Pasteczka)

For a family of continuous functions $f_1, f_2, \dots : I \rightarrow \mathbb{R}$ (I is a fixed closed interval) with $f_1 \leq f_2 \leq \dots$ define a set

$$I_f := \left\{ x \in I : \lim_{n \rightarrow \infty} f_n(x) = +\infty \right\}.$$

We study the properties of the family of all admissible I_f -s and the family of all admissible I_f -s under the additional assumption

$$\lim_{n \rightarrow \infty} \int_x^y f_n(t) dt = +\infty \quad \text{for all } x, y \in I \text{ with } x < y.$$

The origin of this problem is the limit behaviour of quasiarithmetic means.

References

- [1] Gryszka, Karol, and Paweł Pasteczka. "On L^1 and pointwise divergence of continuous functions." *arXiv* (2020): arxiv.org/pdf/2007.12752.pdf.

Polynomial identities satisfied by generalized polynomials

The study of additive mappings from a ring into another ring which preserve squares was initiated by G. Ancochea in [1] in connection with problems arising in projective geometry. Later, these results were again strengthened by (among others) Kaplansky [4] and Jacobson–Rickart [3]. It was G. Ancochea who firstly dealt with the connection of Jordan homomorphisms and homomorphisms, see [1] and its related results [3], [4], [5]. Roughly speaking, all the investigated problems were of the following form: Let R and R' be (commutative) rings and $P \in R[x]$ and $Q \in R'[x]$ be polynomials. Determine all those additive functions that also satisfy

$$f(P(x)) = Q(g(x))$$

for all $x \in \mathbb{R}$. Clearly, this problem is meaningful if instead of additive functions, generalized monomials are considered. According to this remark, the talk will be devoted to the following problem: let n be a positive integer, \mathbb{F} be a field and $P \in \mathbb{F}[x]$ and $Q \in \mathbb{C}[x]$ be polynomials. Determine those generalized polynomials $f, g: \mathbb{F} \rightarrow \mathbb{C}$ of degree n that also fulfill equation

$$f(P(x)) = Q(g(x))$$

for each $x \in \mathbb{F}$. As it turns out, the difficulty of such problems heavily relies on that we consider the above equation for generalized polynomials or for (normal) polynomials. Therefore, firstly we study the connection between these two notions.

References

- [1] Ancochea, Germán. "Le théorème de von Staudt en géométrie projective quaternionnienne." *J. Reine Angew. Math.* 184 (1942): 193-198.
- [2] Herstein, Israel Nathan. "Jordan homomorphisms." *Trans. Amer. Math. Soc.* 81 (1956): 331-341.
- [3] Jacobson, Nathan, and Charles Earl Rickart. "Jordan homomorphisms of rings." *Trans. Amer. Math. Soc.* 69 (1950): 479-502.
- [4] Kaplansky, Irving. "Semi-automorphisms of rings." *Duke Math. J.* 14 (1947): 521-525.
- [5] Żelazko, Wiesław. "A characterization of multiplicative linear functionals in complex Banach algebras." *Studia Math.* 30 (1968): 83-85.

Grzegorz Guzik

AGH University of Science and Technology, Kraków, Poland

On geometric rate of convergence for generalized random iterations (joint work with Rafał Kapica)

Fix $m \in \mathbb{N}$. Let (X, ϱ) be a complete and separable metric space and $(\Omega, \Sigma, \mathbb{P})$ be a probability space. We consider a *generalized iterated function system with constant probabilities* (GIFSP), i.e. \mathbb{P} -continuous family $\mathcal{F} = \{f_\omega: X^m \rightarrow X : \omega \in \Omega\}$.

Given an infinite word $\omega^\infty = (\omega_n)_{n \in \mathbb{N}} \in \Omega^\infty$ we define a (random) ω^∞ -trajectory of a GIFSP \mathcal{F} starting from $x_1, \dots, x_m \in X$ as a sequence $(x_n)_{n \in \mathbb{N}}$ of points from X defined by

$$x_{n+m} = f_{\omega_n}(x_n, \dots, x_{n+m-1}) \quad \text{for } n \in \mathbb{N}. \quad (1)$$

The above formula is a generalization of well known iteration process for so-called random valued functions studied by many authors on the ground of functional equations as well as operator theory. We study the long-time behavior of random trajectories (1) of GIFSP with arbitrary number of transformations. To do this we discover the theory of limiting behavior for the pretty large class of operators of Markov-type acting on the product of spaces of Borel measures on X . We show the simple criterion on the existence of the unique invariant measure for such operator (asymptotic stability). Moreover, we prove that this unique measure is a probability one with the first moment finite, it attracts all trajectories of distributions under the action of the operator in the sense of the Hutchinson–Wasserstein norm as well in the weak topology. The rate of convergence is the geometric one.

One can apply this result for studying asymptotic behavior of solutions of stochastic difference equations with multiple delays, for example in the case when such equation is a discretization of stochastic differential Itô one.

References

- [1] Guzik, Grzegorz, and Rafał Kapica. "On the geometric ergodicity for a generalized IFS with probabilities". *Stoch. Dyn.*, to appear.

László Horváth

University of Pannonia, Hungary

Refinements of the integral Jensen's inequality generated by finite or infinite permutations

There are a lot of papers dealing with applications of the so called cyclic refinement of the discrete Jensen's inequality. A significant generalization of the cyclic refinement, based on combinatorial considerations, has recently been discovered. In the present talk we give the integral versions of these results. On the one hand, a new method to refine the integral Jensen's inequality is developed. Furthermore, the result contains some recent refinements of the integral Jensen's inequality as elementary cases. Finally, some applications to the Fejér inequality (especially the Hermite-Hadamard inequality), quasi-arithmetic means and f -divergences are presented.

Eliza Jabłońska

AGH University of Science and Technology, Kraków, Poland

Remarks on classes $\mathcal{A}, \mathcal{B}, \mathcal{C}$

Ger and Kuczma [4] introduced the following three families of sets in X :

- $\mathcal{A}(X)$ of all subsets $T \subset X$ such that any mid-convex function $f: D \rightarrow \mathbb{R}$ defined on a convex open subset $D \subset X$ containing T is continuous if $\sup f(T) < \infty$;
- $\mathcal{B}(X)$ of all subsets $T \subset X$ such that any additive function $f: X \rightarrow \mathbb{R}$ with $\sup f(T) < \infty$ is continuous;
- $\mathcal{C}(X)$ of all subsets $T \subset X$ such that any additive function $f: X \rightarrow \mathbb{R}$ is continuous if the set $f(T)$ is bounded in \mathbb{R} .

It is clear that

$$\mathcal{A}(X) \subset \mathcal{B}(X) \subset \mathcal{C}(X). \quad (1)$$

By the example of Erdős [2] (discussed in [4]) the classes $\mathcal{B}(X)$ and $\mathcal{C}(X)$ are not equal even if $X = \mathbb{R}^n$, $n \in \mathbb{N}$. On the other hand, Ger and Kominek [3] proved that $\mathcal{A}(X) = \mathcal{B}(X)$ for any Baire topological vector space X . In particular, $\mathcal{A}(\mathbb{R}^n) = \mathcal{B}(\mathbb{R}^n)$ for every $n \in \mathbb{N}$ (cf. [5]).

There are lots of papers devoted to the problem of recognizing sets in the families $\mathcal{A}(X)$, $\mathcal{B}(X)$ or $\mathcal{C}(X)$. Some recent results in [1] allow us to give some further examples of sets which are (not) in $\mathcal{A}(X)$, $\mathcal{B}(X)$ or $\mathcal{C}(X)$.

References

- [1] Banach, Taras, Szymon Głąb, Eliza Jabłońska, and Jarosław Swaczyna. "Haar- \mathcal{I} sets: looking at small sets in Polish groups through compact glasses." *Dissertationes Math.* (2021). <https://doi.org/10.4064/dm812-2-2021>
- [2] Erdős, Paul. "On some properties of Hamel bases." *Colloq. Math.* 10 (1963): 267-269.
- [3] Ger, Roman, and Zygfryd Kominek. "Boundedness and continuity of additive and convex functionals." *Aequationes Math.* 37, no. 2-3 (1989): 252-258.
- [4] Ger, Roman. "On the boundedness and continuity of convex functions and additive functions." *Aequationes Math.* 4 (1970): 157-162.
- [5] Kuczma, Marcin Emil. "On discontinuous additive functions." *Fund. Math.* 66 (1969/70): 383-392.

Justyna Jarczyk

University of Zielona Góra, Poland

On generalized Archimedes–Borchardt algorithm (joint work with Witold Jarczyk)

We investigate convergence and invariance properties of the generalized Archimedes–Borchardt algorithm. The main tool is reducing the problem to an appropriate Gauss iteration process.

Witold Jarczyk
University of Zielona Góra, Poland

Form of semiflows of pairs of weighted quasi-arithmetic means
(joint work with Dorota Głazowska, Justyna Jarczyk and Janusz Matkowski)

We start with a reformulation of Theorem 4 from the paper [1], more friend for our consecutive considerations than the original one. It provides a clear description of a general form of semiflows of pairs of weighted quasi-arithmetic means, with time running through the set \mathbb{D}_+ of all positive dyadic numbers (see Theorem 1). Because of the density of \mathbb{D}_+ in $(0, +\infty)$ we obtain the form of all continuous semiflows of such a type, over $(0, +\infty)$ (see Corollary 2).

In the second part of the talk we present an equivalent version of Theorem 1 from the paper [2] and observe that this statement runs contrary to Corollary 2. The reasons of such a situation are presented.

Finally we ask if the assumption of the continuity of semiflow is essential for the validity of Corollary 2.

References

- [1] Głazowska, Dorota, Justyna Jarczyk, and Witold Jarczyk. "Embeddability of pairs of weighted quasi-arithmetic means into a semiflow." *Aequationes Math.* 94, no. 4 (2020): 679-687.
Correction: *Aequationes Math.* 95, no. 3 (2021): 599-600.
- [2] Matkowski, Janusz. "On iteration semigroups of mean-type mappings and invariant means." *Aequationes Math.* 64, no. 3 (2002): 297-303.

Sándor Jenei
University of Pécs, Hungary

Algebraic methods in functional equations – beyond the regularity conditions

In [2, 3] it has been shown how algebraic and geometric methods can be used to solve an open problem posed in [1]. We shall shortly describe an example which demonstrates that when considering associative functions, algebraic methods can go much beyond even the most general regularity condition which is assumed in functional equations, namely the continuity of the solution, and one can still obtain all solutions. Moreover, the domain of the functions in our example is not necessarily a real interval either, it can be any linearly (totally) ordered set. The applied techniques (the theory of residuated semigroups) often work on lattices and even on partially ordered sets. Under the following conditions we describe all solutions of the associativity equation

$$F(x, F(y, z)) = F(F(x, y), z)$$

- for $x, y \in X$ such that $F(x, y) = F(y, x)$,
- the domain of the function F is $X \times X$, X is any linearly ordered set,
- there exists $t \in X$ such that $F(x, t) = x$,
- for any $x, y \in X$ there exists the greatest element $F_{\rightarrow}(x, y)$ of the set $\{z \in X : F(x, z) \leq y\}$
- for $x \in X$, $F_{\rightarrow}(F_{\rightarrow}(x, t), t) = x$,
- $F_{\rightarrow}(t, t) = t$.

Or using an infix notation $x \circledast (y \circledast z) = (x \circledast y) \circledast z$, where

- \circledast is a binary operation over a linearly ordered set X ,
- for $x, y \in X$, $x \circledast y = y \circledast x$,
- there exists $t \in X$ such that $x \circledast t = x$,
- for any $x, y \in X$ there exists the greatest element $x \rightarrow_{\circledast} y$ of the set $\{z \in X \mid x \circledast z \leq y\}$,
- for $x \in X$, $(x \rightarrow_{\circledast} t) \rightarrow_{\circledast} t = x$,
- $t \rightarrow_{\circledast} t = t$

The fourth items say that F (resp. \circledast) is residuated. It implies that \circledast is monotone increasing, and hence it can be also regarded as left-continuity.

This talk is based on the article [4].

References

- [1] Alsina, Claudi, Maurice Jerome Frank, and Berthold Schweizer. "Problems on associative functions." *Aequationes Math.* 66, no. 1-2 (2003): 128-140.
- [2] Jenei, Sándor. "On the convex combination of left-continuous t -norms." *Aequationes Math.* 72, no. 1-2 (2006): 47-59.
- [3] Jenei, Sándor. "On the Geometry of Associativity." In: *Report of Meeting "15th International Conference on Functional Equations and Inequalities"*, Ustroń, May 19–25, 2013. Ann. Univ. Paedagog. Crac. Stud. Math. 12 (2013), 91–131.
- [4] Jenei, Sándor. "Group representation for even and odd involutive commutative residuated chains." *arXiv* (2019): arxiv.org/pdf/1910.01404.pdf.

Soon-Mo Jung

Hongik University, Seoul, Republic of Korea

Approximate isometries on bounded sets

More than 20 years after Fickett [2] attempted to prove the Hyers-Ulam stability of isometries defined on a bounded subset of \mathbb{R}^n in 1982, Alestalo et al. [1] and Väisälä [4] improved the Fickett's theorem significantly. In this lecture, using a more intuitive and efficient approach, first used in the paper [3], we improve Fickett's theorem by proving the Hyers-Ulam stability of isometries defined on a bounded subset of \mathbb{R}^n .

References

- [1] Alestalo, Pekka, Dmitry Alexandrovich Trotsenko, and Jussi Väisälä. "Isometric approximation." *Israel J. Math.* 125 (2001): 61-82.
- [2] Fickett, James Wildon. "Approximate isometries on bounded sets with an application to measure theory." *Studia Math.* 72, no. 1 (1982): 37-46.
- [3] Jung, Soon-Mo. "Hyers-Ulam stability of isometries on bounded domains." *Open Math.* 19, no. 1 (2021): 675-689.
- [4] Väisälä, Jussi. "Isometric approximation property in Euclidean spaces." *Israel J. Math.* 128 (2002): 1-27.

Gergely Kiss

Alfréd Rényi Institute of Mathematics, Budapest, Hungary

Charaterization of quasi-arithmetic means without regularity (joint work with Pál Burai and Patrícia Szokol)

In this talk we recall the classical result of Aczél for the characterization of quasi-arithmetic means. We analysis the regularity assumption of this construction and we show that every bisymmetric, partially strictly monotonic, reflexive and symmetric function $F: I^2 \rightarrow I$ is continuous, where I is a proper subinterval of \mathbb{R} . This talk is based on the article [1] and the book [2].

References

- [1] Burai, Pál, Gergely Kiss, and Patrícia Szokol. "Characterization of quasi-arithmetic means without regularity condition". Submitted to *Acta Math. Hungar.*. *arXiv* (2021): arxiv.org/pdf/2107.07391.pdf.
- [2] Aczél, János, and Jean Dhombres. *Functional equations in several variables*. Vol. 31 of *Encyclopedia of Mathematics and its Applications*. Cambridge: Cambridge University Press, 1989.

Grzegorz Kleszcz

AGH University of Science and Technology, Kraków, Poland

On a functional–difference inclusion induced by some iterated function system (joint work with Grzegorz Guzik)

In [1, Part II, Chapter 1] the following functional–difference equations with continuous–time argument

$$x(t+1) = f(x(t)) \quad (1)$$

is considered. In general, here above f is a given continuous function of some compact space X into itself and $x: [0, \infty) \rightarrow X$ is unknown. One can see that solutions of (1) depend on an arbitrary function. More precisely, having a function $\varphi: [0, 1] \rightarrow X$ and putting

$$x|_{[0,1]} := \varphi \quad (2)$$

one can get uniquely the solution x on the whole semiline $[0, \infty)$ by the formula

$$x(t+n) = f^n \circ \varphi(t) \quad \text{for } n \in \mathbb{N}. \quad (3)$$

In particular, if φ is a continuous function and the initial condition

$$\varphi(1) = f(\varphi(0)). \quad (4)$$

is assumed, the solution x of (1) is continuous.

The most interesting aspect is a study of asymptotic behaviors of graphs of a sequence $(f^n \circ \varphi)$ of compositions of iterates f^n with an 'observable' function φ in a proper Hausdorff metric or the topology of uniform convergence. In many cases such graphs tend to the graph of some set–valued function (with closed graph) rather than to the graph of a single–valued continuous function. Moreover, it is remarkable that usually the asymptotic behavior of such a sequence of graphs strongly depends on a choice of a function φ .

In various models of real systems the given function is not uniquely determined but either can be chosen from some family (finite or not) by a deterministic or random algorithm, or it is determined with some error. Hence it seems quite reasonable to consider the following inclusion

$$x(t+1) \in f(x(t)) \quad (5)$$

instead of the equality (1). Here $f: X \rightsquigarrow X$ is a given set–valued function which can represent a union of all functions possibly chosen.

In the present talk we consider the inclusion (5) with set–valued function f generated by an iterated function system, i.e. a finite family of continuous mappings of X into itself, which is assumed to possess an attractor (a fractal set). Our main result says that under quite typical assumptions on a given iterated function system we get a simple asymptotic behavior of graphs of compositions $f^n \circ \varphi$, $n \in \mathbb{N}$ which is, surprisingly, independent on a choice of an 'observable' φ .

References

- [1] Sharkovsky, Oleksandr M., Yury L. Maistrienko, and Elena Yu. Romanenko. *Difference equations and their applications*. Vol. 250 of *Mathematics and Its Applications*. New York: Springer-Verlag, 1993.

Approximately order zero maps between C^* -algebras

A bounded linear operator $\phi: A \rightarrow B$ acting between two C^* -algebras is called an ε -order zero map, provided that $\|\phi(x)\phi(y)\| \leq \varepsilon\|x\|\|y\|$ for all positive $x, y \in A$ with $xy = 0$. This terminology arises naturally from the theory of order zero maps and nuclear dimension of C^* -algebras developed by Winter and Zacharias in [2] and [3]. In that theory, completely positive order zero maps (i.e. preserving zero products of positive elements) correspond to usual partitions of unity in the commutative setting, and they serve as building blocks for the notion of nuclear dimension which is a noncommutative analogue of the topological covering dimension.

During the talk, we will report on selected results from [1] which yield some structural properties of ε -order zero maps concerning approximate Jordan-like equations and almost commutation relations. The main goal is to decide, for a given pair (A, B) of C^* -algebras, whether and under what conditions any ε -order zero map $\phi: A \rightarrow B$ can be approximated by an approximate Jordan $*$ -homomorphism, with both errors of approximation depending only on $\|\phi\|$ and ε . As we shall explain, this is for instance possible in the case where the codomain algebra B is finite-dimensional. In particular, if $B = \mathbb{M}_n(\mathbb{C})$ is the $n \times n$ matrix algebra, then for an arbitrary C^* -algebra A , any $0 < \varepsilon \leq 1$, and any positive ε -order zero map $\phi: A \rightarrow \mathbb{M}_n(\mathbb{C})$, there exist a corner C^* -subalgebra $C \subseteq \mathbb{M}_n(\mathbb{C})$ and a bounded linear operator $\Phi: A_1 \rightarrow C$ defined on the unitization of A satisfying

$$\|\phi - \Phi\| \leq 37\|\phi\|^{4/5}\varepsilon^{1/16}$$

and such that either $\Phi = 0$ or $\Phi(1)$ is invertible in C in which case the operator $\Phi(1)^{-1}\Phi(\cdot)$ is an approximate Jordan $*$ -homomorphism with an error δ estimated by

$$\delta = \mathcal{O}(256^n)\|\phi\|\varepsilon^{1/16}.$$

A similar result is valid in the case where A is commutative, B is arbitrary and $\phi: A \rightarrow B$ is assumed to be a surjective self-adjoint ε -order zero map. In this situation however, an approximating operator takes values in a corner subalgebra of the bicommutant B'' .

References

- [1] Kochanek, Tomasz. "Approximately order zero maps between C^* -algebras." *J. Funct. Anal.* 281, no. 2 (2021): Paper No. 109025.
- [2] Winter, Wilhelm, and Joachim Zacharias. "Completely positive maps of order zero." *Münster J. Math.* 2 (2009): 311-324.
- [3] Winter, Wilhelm, and Joachim Zacharias. "The nuclear dimension of C^* -algebras." *Adv. Math.* 224, no. 2 (2010): 461-498.

Zbigniew Leśniak 
Pedagogical University of Krakow, Poland

On a construction of iterative roots of a Brouwer homeomorphism

We present a method of construction of continuous orientation preserving iterative roots of a Brouwer homeomorphism under the assumption that for this homeomorphism there exists a family of pairwise disjoint invariant lines covering the plane which satisfies a matching property. The roots are constructed recursively for a family of maximal parallelizable regions, each of which is the union of the invariant lines. The matching property is used to obtain the continuity of the constructed iterative roots.

References

- [1] Le Roux, Frédéric, Anthony G. O'Farrell, Maria Roginskaya, and Ian Short. "Flowability of plane homeomorphisms." *Ann. Inst. Fourier (Grenoble)* 62, no. 2 (2012): 619-639.
- [2] Leśniak, Zbigniew. "On fractional iterates of a Brouwer homeomorphism embeddable in a flow." *J. Math. Anal. Appl.* 366, no. 1 (2010): 310-318.

Radosław Łukasik

University of Silesia, Katowice, Poland

Vector-valued invariant means and projections from the bidual space

Invariant means on amenable groups are an important tool in many parts of mathematics, especially in harmonic analysis. Invariant means and their generalizations for vector-valued functions play also an important role in the stability of functional equations and selections of set-valued functions. Some generalization of the invariant mean for vector-valued functions was investigated in [2] and the existence of such means is connected with reflexive spaces.

Some generalized definition of an invariant mean has been used by many mathematicians as a folklore (e.g. by A. Pełczyński [5]). The explicit form of this definition we can find, e.g. in the work of R. Ger [3]. The space of all bounded functions from a set S into a Banach space X is denoted by $\ell_\infty(S, X)$.

Definition. Let $(S, +)$ be a left [right] amenable semigroup, X be a real Banach space. A linear map $M: \ell_\infty(S, X) \rightarrow X$ is called left [right] X -valued invariant mean if

$$\begin{aligned} \|M\| &\leq 1, \\ M(c\mathbb{1}_S) &= c, \quad c \in X, \\ M({}_a f) &= M(f), \quad a \in S, f \in \ell_\infty(S, X), \\ [M(f_a) &= M(f), \quad a \in S, f \in \ell_\infty(S, X),] \end{aligned}$$

where

$$\begin{aligned} {}_a f(x) &= f(a + x), \quad a, x \in S, \quad f \in \ell_\infty(S, X), \\ [f_a(x) &= f(x + a), \quad a, x \in S, \quad f \in \ell_\infty(S, X).] \end{aligned}$$

If M is a left and right invariant mean, then M is called X -valued invariant mean.


If in the above definition norm of map M is equal at most $\lambda \geq 1$, then M is called X -valued invariant λ -mean.

The existence of such invariant means for a fixed Banach space and for all amenable semigroups has been studied by H.B. Domecq [1, Theorem 2] (his paper has a gap in the proof which was corrected by T. Kania [4]).

In this talk we will show a connection between the existence of X -valued invariant λ -means on X and projections from subspaces of bidual space X^{**} (with large enough density) onto X .

References

- [1] Bustos Domecq, H. "Vector-valued invariant means revisited." *J. Math. Anal. Appl.* 275, no. 2 (2002): 512-520.
- [2] Gajda, Zbigniew. *Invariant means and representations of semigroups in the theory of functional equations*. Katowice: Prace Naukowe Uniwersytetu Śląskiego, 1992.
- [3] Ger, Roman. "The singular case in the stability behavior of linear mappings." *Grazer Math. Ber.* 316 (1992): 59-70.
- [4] Kania, Tomasz. "Vector-valued invariant means revisited once again." *J. Math. Anal. Appl.* 445, no. 1 (2017): 797-802.
- [5] Pełczyński, Aleksander. "Linear extensions, linear averagings, and their applications to linear topological classification of spaces of continuous functions." *Dissertationes Math. (Rozprawy Mat.)* 58 (1968): 92 pp.

Arpita Mal 

Jadavpur University, Kolkata, India

Extreme contraction from the perspective of multi-smoothness

In this talk we explore a relation between extreme contraction and multi-smoothness of bounded linear operators on polyhedral Banach spaces. Using this relation we describe extreme contractions on two-dimensional polygonal Banach spaces. We provide a large class of Banach spaces \mathbb{X}, \mathbb{Y} such that for each extreme contraction $T : \mathbb{X} \rightarrow \mathbb{Y}$ there is an extreme point $x \in \mathbb{X}$ such that Tx is an extreme point of \mathbb{Y} . We also show how the discussed approach helps us to explicitly compute the number of extreme contractions on some Banach spaces. This talk is based on the paper [1].

References

- [1] Mal, Arpita, Kallol Paul, and Subhrajit Dey. "Characterization of extreme contractions through k -smoothness of operators." *Linear Multilinear Algebra* (2021). <https://doi.org/10.1080/03081087.2021.1913086>

Daniela Marian 

Technical University of Cluj-Napoca, Romania

On Hyers-Ulam-Rassias stability of certain partial differential equations and integral equations

In the first part we will present the Ulam-Hyers stability and Ulam-Hyers-Rassias stability for the Darboux-Ionescu problem. This part is based on the article [1] .

In the second part we will present some results for a Volterra-Hammerstein integral equation with modified arguments: existence and uniqueness, integral inequalities, monotony and Ulam-Hyers-Rassias stability. This part is based on the article [2].

References

- [1] Marian, Daniela, Sorina Anamaria Ciplea, and Nicolaie Lungu. "On a functional integral equation". *Symmetry* 13, no. 8 (2021): Article no. 1321.
- [2] Marian, Daniela, Sorina Anamaria Ciplea, and Nicolaie Lungu. "Ulam-Hyers stability of Darboux-Ionescu problem." *Carpathian J. Math.* 37, no. 2 (2021): 211-216.

Joanna Markowicz

Pedagogical University of Krakow, Poland

Uniform convexity of direct sums and interpolation spaces

A natural question in the theory of direct sums and interpolation spaces is whether given property of Banach spaces can be preserved under passing to a direct sum or an interpolation space.

In this talk, we will recall the well known theory of uniform convexity.

Subsequently, we will present the estimation of the modulus of convexity and characteristic of convexity of a general direct sum of Banach spaces.

Next, interpolation spaces obtained with the use of the general discrete interpolation method based on an abstract space with an unconditional basis will be introduced. We will give also conditions which guarantee such interpolation spaces be uniformly convex.

References

- [1] Markowicz, Joanna, and Stanisław Prus. "Uniform convexity of general direct sums and interpolation spaces." *J. Topol. Anal.* (2021): to appear.

Janusz Matkowski

University of Zielona Góra, Poland

Remarks on fixed point theorem

A generalization of Browder-Göhde-Kirk fixed point theorem and its application will be presented.

Rayene Menzer 

University of Debrecen, Hungary

Alternative equations for quadratic functions (joint work with Zoltán Boros)

Kominek, Reich and Schwaiger [2] investigated additive functions that satisfy the additional equation

$$f(x)f(y) = 0 \tag{1}$$

for every $(x, y) \in D$, considering various subsets D of \mathbb{R}^2 , involving, for instance, the graphs of polynomials and the unit circle.

The result for the unit circle was extended to the case when f is a generalized polynomial by Boros and Fechner [1]. On the other hand, Kutas [3] proved the existence of a not identically zero additive function f that satisfies (1) whenever $xy = 1$.

Motivated by these preliminaries, we investigate quadratic functions f that satisfy (1) for every $(x, y) \in D$, considering various subsets D of \mathbb{R}^2 . For the case when D is the graph of a polynomial, we establish the following statement.

Theorem 1. *If $n \in \mathbb{N}$, $a_j \in \mathbb{R}$ ($j = 0, 1, \dots, n-1$) and the quadratic function $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies the equation*

$$f(x) \cdot f\left(x^n + \sum_{k=0}^{n-1} a_k x^k\right) = 0$$

for every $x \in \mathbb{R}$, then $f(x) = 0$ identically.

While Kutas' counterexample for the hyperbola $xy = 1$ can be easily extended to quadratic functions, we can establish a positive result for another hyperbola.

Theorem 2. *If $f: \mathbb{R} \rightarrow \mathbb{R}$ is a quadratic function fulfilling $f(x)f(y) = 0$ whenever $x^2 - y^2 = 1$, then $f(x) = 0$ identically for every $x \in \mathbb{R}$.*

References

- [1] Boros, Zoltán, and Włodzimierz Fechner. "An alternative equation for polynomial functions." *Aequationes Math.* 89, no. 1 (2015): 17-22.
- [2] Kominek, Zygfryd, Ludwig Reich, and Jens Schwaiger. "On additive functions fulfilling some additional condition." *Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II* 207 (1998): 35-42 (1999).
- [3] Kutas, Péter. "Algebraic conditions for additive functions over the reals and over finite fields." *Aequationes Math.* 92, no. 3 (2018): 563-575.

Flavia-Corina Mitroi-Symeonidis
Academy of Economic Studies, Bucharest, Romania

Redistributing algorithms and Shannon's Entropy
(joint work with Eleutherius Symeonidis)

The mathematical problems discussed in our talk arise in the framework of time series analysis, namely in the analysis of temperature values measured during compartment fire experiments. The permutation entropy is used for such analysis, and the underlying permutations were established using redistributing algorithms. We investigate how Shannon's entropy changes when the distribution probability is modified by means of redistributing algorithms.

Let a probability distribution $P = (p_1, \dots, p_n)$ be such that it has at least k nonzero components p_{n-k+1}, \dots, p_n ($1 \leq k \leq n-1$). We are interested to find the best (largest) domain $\Delta \subset \mathbb{R}^k$ such that

$$H(p_1, \dots, p_{n-k}, p_{n-k+1}, \dots, p_n) \geq H(0, \dots, 0, 1/k, \dots, 1/k)$$

for all P with $(p_{n-k+1}, \dots, p_n) \in \Delta$.

We mention the following immediate consequences of interest, which we assume that are already known (we have not found any reference for them yet):

Let n , k and p be fixed. The entropy $H(p_1, \dots, p_{n-k}, p, \dots, p)$ attains its maximum at

$$p_1 = \dots = p_{n-k-1} = p_{n-k} = \frac{1 - kp}{n - k},$$

and its minimum if there is $j = 1, \dots, n - k$ such that $p_j = 1 - kp$.

Mădălina Moga

Babeş-Bolyai University, Cluj-Napoca, Romania

On some properties of Meir-Keeler operators

The first purpose of this talk is to prove that, in a Banach space X , a Meir-Keeler operator f (singlevalued or multivalued) is a norm-contraction. Then, using this result, we will give sufficient conditions assuring that the field $1_X - f$, generated by f , is surjective. It also contains surjectivity theorems for singlevalued and, respectively, multivalued Meir-Keeler operators. In the last part we present an application for the previous results. Our results generalize some well-known theorems of this type for Banach/Nadler type contractions, see [5] and [2], as well as other results of this type for generalized contractions, see also [1], [3], [6], [7].

We will present a new surjectivity theorem for a singlevalued Meir-Keeler operator. The approach is based on the norm-contraction operator theory and on the well-known theorem of A. Granas [4].

Theorem 1. *Let $(X, \|\cdot\|)$ be a Banach space and Y be a nonempty, closed and convex subset of X . Let us consider the operator $f: Y \rightarrow Y$ satisfying the following assumptions:*

1. *f restricted to any bounded set in Y is compact;*
2. *f is a Meir-Keeler operator.*

Then $1_Y - f$ is a surjective operator.

Our main result concerning the surjectivity of the multivalued field generated by a Meir-Keeler operator is the following theorem.

Theorem 2. *Let $(X, \|\cdot\|)$ be a Banach space and Y be a nonempty, closed and convex subset of X . Let us consider the multivalued operator $T: Y \rightarrow P_{cp,cv}(Y)$ satisfying the following assumptions:*

1. *for each $A \in P_b(X)$ the set $T(A)$ is relatively compact;*
2. *T is a Meir-Keeler operator.*

Then, the field $1_Y - T$ generated by T is surjective.

References

- [1] Aldea, Florica. "Surjectivity theorems for normcontraction operators." *Mathematica* 44(67) (2003), no. 2: 129-136.
- [2] Andres, Jan, and Lech Górniewicz. "On the Banach contraction principle for multivalued mappings." In: M. Lassonde (ed.) *Approximation, Optimization and Mathematical Economics*. Heidelberg: Physica, 2001, 1-23.
- [3] Anisiu, Mira-Cristiana. "Quasibounded mappings and generalized contraction." Seminar on Fixed Point Theory Cluj-Napoca, Preprint, 83-3, Univ. "Babeş-Bolyai", Cluj-Napoca (1983): 131-134.
- [4] Granas, Andrzej. "On a certain class of nonlinear mappings in Banach space." *Bull. Acad. Pol. Sci.* 9 (1957): 867-871.
- [5] Granas, Andrzej, and James Dugundji. *Fixed point theory*. Springer Monographs in Mathematics. New York: Springer-Verlag, 2003.
- [6] Petruşel, Gabriela. "Generalized multivalued contractions which are quasi-bounded." *Demonstratio Math.* 40, no. 3 (2007): 639-648.
- [7] Rus, Ioan A. "Normcontraction mappings outside a bounded set". *Itinerant Seminar on Functional Equations Approximation and Convexity*, Cluj-Napoca (1986), 257-260.

Lajos Molnár 

University of Szeged, Hungary

Budapest University of Technology and Economics, Hungary

On the order determining property of the norm of a Kubo-Ando mean in operator algebras

Recently, several papers have been published concerning preservers of norms of means on function algebras and operator algebras. In our approach to the study of those transformations and in the proofs of our results the key point was an order determining property of the norms of the considered means.

Motivated by that, in a current paper we have initiated the investigation of the mentioned property in a general setting. The aim of this talk is to report on the obtained results concerning the question of when the operator norm of a Kubo-Ando mean σ determines the order on the positive definite cone of an operator algebra. More precisely, the question asks when, for any given pair A, B of positive definite elements of the underlying algebra, we have that $A \leq B$ holds if and only if $\|A\sigma X\| \leq \|B\sigma X\|$ is valid for all positive definite elements X . Our study is not completed, besides presenting results we formulate open problems.

References

- [1] Molnár, Lajos. "On the order determining property of the norm of a Kubo-Ando mean in operator algebras." *Integral Equations and Operator Theory* 93 (2021): Article No. 53.

Veerapazham Murugan 

National Institute of Technology Karnataka, Mangaluru, India

Iterative root problem for continuous functions of non-monotonicity height 1

In this paper, we define the non-monotonicity height for any continuous self-maps on a compact interval and prove some interesting properties of the non-monotonicity height of non-PM functions. We obtain necessary and sufficient conditions for the existence of iterative roots for a class of continuous functions of non-monotonicity height 1.

Fixed point results on relational metric spaces with applications to Hyers-Ulam stability and nonlinear matrix equations

Throughout this talk, we will develop some fixed point results for \mathcal{FG} -contractive mappings on complete metric spaces equipped with any binary relation (not necessarily a partial order). Additionally, the Ulam-Hyers stability, well-posedness, and limit shadowing properties of this issue are examined. We use the aforementioned fixed point result to find the solution to a non-linear matrix equation of the form $\mathcal{X} = \mathcal{Q} + \sum_{i=1}^m \mathcal{A}_i^* \mathcal{G}(\mathcal{X}) \mathcal{A}_i$, where \mathcal{Q} is a Hermitian positive definite matrix, \mathcal{A}_i^* stands for the conjugate transpose of an $n \times n$ matrix \mathcal{A}_i and \mathcal{G} is an order-preserving continuous mapping from the set of all Hermitian matrices to the set of all positive definite matrices such that $\mathcal{G}(O) = O$. We explore the necessary and sufficient criteria for the existence of a unique positive definite solution to a particular matrix problem. Through several demonstrations using graphical representations, we demonstrate the fixed-point conclusions and the relevance of related work, as well as the convergence analysis of non-linear matrix equations.

Definition . [1] The collection of all functions $\mathcal{F}: \mathbb{R}_+ \rightarrow \mathbb{R}$ satisfying:

(\mathcal{F}_1) \mathcal{F} is continuous and strictly increasing;

(\mathcal{F}_2) for each $\{\xi_n\} \subseteq \mathbb{R}_+$, $\lim_{n \rightarrow \infty} \xi_n = 0$ iff $\lim_{n \rightarrow \infty} \mathcal{F}(\xi_n) = -\infty$,

will be denoted by \mathbb{F} .

The collection of all pairs of mappings (\mathcal{G}, β) , where $\mathcal{G}: \mathbb{R}_+ \rightarrow \mathbb{R}$, $\beta: \mathbb{R}_+ \rightarrow [0, 1)$, satisfying:

(\mathcal{F}_3) for each $\{\xi_n\} \subseteq \mathbb{R}_+$, $\limsup_{n \rightarrow \infty} \mathcal{G}(\xi_n) \geq 0$ iff $\limsup_{n \rightarrow \infty} \xi_n \geq 1$;

(\mathcal{F}_4) for each $\{\xi_n\} \subseteq \mathbb{R}_+$, $\limsup_{n \rightarrow \infty} \beta(\xi_n) = 1$ implies $\lim_{n \rightarrow \infty} \xi_n = 0$;

(\mathcal{F}_5) for each $\{\xi_n\} \subseteq \mathbb{R}_+$, $\sum_{n=1}^{\infty} \mathcal{G}(\beta(\xi_n)) = -\infty$,

will be denoted by \mathbb{G}_β .

We call $(\mathcal{X}, \mathfrak{R})$ a relational set if (i) $\mathcal{X} \neq \emptyset$ is a set and (ii) \mathfrak{R} is a binary relation on \mathcal{X} .

In addition, if (\mathcal{X}, d) is a metric space, we call $(\mathcal{X}, d, \mathfrak{R})$ a relational metric space (*RMS*, for short).

Definition . Let $(\mathcal{X}, d, \mathfrak{R})$. A mapping *RMS* and $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$ is said to be a \mathcal{FG} -contractive mapping, if there are $\mathcal{F} \in \mathbb{F}$ and $(\mathcal{G}, \beta) \in \mathbb{G}_\beta$, such that for $(\nu, \vartheta) \in \mathcal{X}$ with $(\nu, \vartheta) \in \mathfrak{R}^*$,

$$\mathcal{F}(d(\mathcal{T}\nu, \mathcal{T}\vartheta)) \leq \mathcal{F}(\Theta(\nu, \vartheta)) + \mathcal{G}(\beta(\Theta(\nu, \vartheta))),$$

where $\Theta(\nu, \vartheta) = \max \left\{ d(\nu, \vartheta), d(\nu, \mathcal{T}\nu), d(\vartheta, \mathcal{T}\vartheta), \frac{d(\nu, \mathcal{T}\vartheta) + d(\vartheta, \mathcal{T}\nu)}{2} \right\}$ and $\mathfrak{R}^* = \{(\nu, \vartheta) \in \mathfrak{R} \mid \mathcal{T}\nu \neq \mathcal{T}\vartheta\}$. We denote by $(FG)_{\mathfrak{R}}$ the collection of all \mathcal{FG} -contractive mappings on $(\mathcal{X}, d, \mathfrak{R})$.

Theorem 1. Let $(\mathcal{X}, d, \mathfrak{R})$ be an *RMS* and $\mathcal{T}: \mathcal{X} \rightarrow \mathcal{X}$. Suppose that the following conditions hold: (i) $\mathfrak{X}(\mathcal{T}, \mathfrak{R}) \neq \emptyset$; (ii) \mathfrak{R} is \mathcal{T} -closed and \mathcal{T} -transitive; (iii) \mathcal{X} is \mathcal{T} - \mathfrak{R} -complete; (iv) $\mathcal{T} \in (FG)_{\mathfrak{R}}$; (v) \mathfrak{S} is \mathfrak{R} -continuous or (vi) \mathfrak{R} is d -self-closed. Then there exists a fixed point $\nu_* \in \mathcal{X}$ of \mathcal{T} .

References

- [1] Parvaneh, Vahid, Nawab Hussain, and Zoran Kadelburg. "Generalized Wardowski type fixed point theorems via α -admissible FG -contractions in b -metric spaces." *Acta Math. Sci. Ser. B (Engl. Ed.)* 36, no. 5 (2016): 1445-1456.

Kazimierz Nikodem
University of Bielsko-Biala, Poland

Remarks on set-valued means

Let X be a real vector space and D be a convex nonempty subset of X . Denote by $S(D)$ the family of all nonempty subsets of D . We say that a function $M: D^n \rightarrow S(D)$ is a *set-valued mean* if

$$M(x_1, \dots, x_n) \subset \text{conv}\{x_1, \dots, x_n\},$$

for all $x_1, \dots, x_n \in D$.

Set-valued counterparts of the arithmetic, quasi-arithmetic and Lagrangian means are investigated and various properties of them are presented. In particular, the following result is presented.

Theorem. *Let $f, g: I \rightarrow J$ be strictly increasing functions such that f is concave and g is convex. Assume that $f \leq g$ on I and $F(x) = [f(x), g(x)]$ holds for all $x \in I$. Then the map $A_F: I^n \rightarrow S(I)$ given by*

$$A_F(x_1, \dots, x_n) = F^+ \left(\frac{1}{n} \sum_{i=1}^n F(x_i) \right), \quad x_1, \dots, x_n \in I,$$

and the map $L_F: I^2 \rightarrow S(I)$ given by

$$L_F(x_1, x_2) = \begin{cases} F^+ \left(\frac{1}{x_2 - x_1} \int_{x_1}^{x_2} F(x) dx \right) & \text{if } x_1 \neq x_2 \\ \{x_1\} & \text{if } x_1 = x_2 \end{cases}$$

are set-valued means.

References

- [1] Nikodem, Kazimierz. "Set-valued means." *Set-Valued Var. Anal.* 28, no. 3 (2020): 559-568.
- [2] Nikodem, Kazimierz. "Characterizations of quasi-affine set-valued means." Submitted.

On some new functional equations characterizing polynomial functions

This is a continuation of the talk presented by Maciej Sablik and containing the theoretical results of the functional equations belonging to the class (1)

$$F(x+y) - F(x) - F(y) = \sum_{i=1}^m (a_i x + b_i y) f(\alpha_i x + \beta_i y) \quad (1)$$

where $F, f : \mathbb{R} \rightarrow \mathbb{R}$ are unknown functions and for each $i \in \{1, \dots, m\}$ the numbers $a_i, b_i \in \mathbb{R}$ and $\alpha_i, \beta_i \in \mathbb{Q}$ are given.

We observed that solving even simple equations belonging to class (1) needs a long and tiresome calculation. Therefore, we developed an algorithm written in the computer algebra system Maple which takes into account the theoretical results to determine the polynomial solutions of the functional equations belonging to the class (1) (cf. also [1], [2] and [3]).

References

- [1] Borus, Gergő Gyula, and Attila Gilányi. "Solving systems of linear functional equations with computer." *2013 IEEE 4th International Conference on Cognitive Infocommunications (CogInfoCom)*, 559-562. Budapest, 2013.
- [2] Borus, Gergő Gyula, and Attila Gilányi. "Computer assisted solution of systems of two variable linear functional equations." *Aequationes Math.* 94, no. 4 (2020): 723-736.
- [3] Gilányi, Attila. "Solving linear functional equations with computer." *Math. Pannon.* 9, no. 1 (1998): 57-70.

Andrzej Olbryś

University of Silesia, Katowice, Poland

On characterization of affine differences and related forms

In the talk we consider the problem of characterizing affine differences and some related expressions. We use functional equations to characterize expressions $\omega: X \times X \times [0, 1] \rightarrow Y$ of the form

$$\omega(x, y, t) = tf(x) + (1 - t)f(y) - f(tx + (1 - t)y), \quad x, y \in X, \quad t \in [0, 1],$$

where $f: X \rightarrow Y$ is an arbitrary function and X, Y are real linear spaces.

References

- [1] Ebanks, Bruce Robertson, and Che Tat Ng. "On generalized cocycle equations." *Aequationes Math.* 46, no. 1-2 (1993): 76-90.
- [2] Ebanks, Bruce Robertson. "Characterization of Jensen differences and related forms." *Aequationes Math.* 76, no. 1-2 (2008): 179-190.
- [3] Erdős, Jenő. "A remark on the paper "On some functional equations" by S. Kurepa." *Glasnik Mat.-Fiz. Astronom. Društvo Mat. Fiz. Hrvatske Ser. II* 14 (1959): 3-5.
- [4] Olbryś, Andrzej. "A sandwich theorem for generalized convexity and applications." *J. Convex Anal.* 26, no. 3 (2019): 887-902.
- [5] Olbryś, Andrzej. "A support theorem for generalized convexity and its applications." *J. Math. Anal. Appl.* 458, no. 2 (2018): 1044-1058.

Adam Ostaszewski

London School of Economics, United Kingdom

Tilted solutions of the Gołab-Schinzel functional equation in Banach algebras (joint work with Nicholas Hugh Bingham)

In \mathbb{R} the Gołab-Schinzel functional equation, briefly: (GS), has continuous solutions affine:

$$S(x) = 1 + \rho x \text{ on } G_S := \{x : 1 + \rho x > 0\} \subseteq \mathbb{R}.$$

Question: For S with values in a Banach algebra \mathbb{A} : what form does a solution S have on

$$\mathbb{G}_S^*(\mathbb{A}) := \{x \in \mathbb{A} : S(x) \in \mathbb{A}^{-1}\}, \text{ or on } \mathbb{G}_S(\mathbb{A}) := \{x \in \mathbb{A} : S(x) \in \mathbb{A}_1\},$$

for \mathbb{A}^{-1} the invertible elements and \mathbb{A}_1 the connected component of the identity?

S is affine iff \mathbb{A} -differentiable (for \mathbb{A}^{-1} dense): compare [2]. Below S is *Fréchet differentiable* with derivative $S'(0)u$ at 0 in direction u . Define the following

$$\begin{aligned} N(x) &: = S(x) - 1_{\mathbb{A}} - \rho x \text{ for } \rho := S(1_{\mathbb{A}}) - 1_{\mathbb{A}} \quad (x \in \mathbb{G}_S^*, \text{ assuming } S(1_{\mathbb{A}}) \in \mathbb{A}^{-1}); \\ \lambda_u(t) &: = (e^{t\gamma(u)} - 1)/(e^{\gamma(u)} - 1) \quad \text{for } \gamma(u) := S'(0)u \text{ with L'Hospital convention}; \\ T(u) &: = u(e^{\gamma(u)} - 1)/\gamma(u) \quad \text{for } u \in \mathbb{A} \text{ (the 'exponential tilting' map)}. \end{aligned}$$

Characterization Theorem (Curvilinear exponential homogeneity). *For S a Fréchet differentiable solution of (GS), N satisfies a (Levi-Civita-style) Goldie equation:*

$$N(x + S(x)y) = N(x) + S(x)N(y) \quad (x, y \in \mathbb{G}_S^*).$$

Furthermore, for all $u \in \mathbb{A}$ with spectrum $\text{spec}(\gamma(u))$ not separating 0 from ∞ :

$$\begin{aligned} N(u(e^{t\gamma(u)} - 1)/\gamma(u)) &= \lambda_u(t)N(u(e^{\gamma(u)} - 1)/\gamma(u)) \quad (t \geq 0), \\ N(tu) &= t1_{\mathbb{A}}N(u) = tN(u) \quad (t \geq 0) \text{ when } 1 \in \exp(\text{spec}(\gamma(u))). \end{aligned}$$

So T exhibits ‘curvilinear exponential homogeneity’ under N : $N(T(tu)) = \lambda_u(t)N(T(u))$.

Corollary. $\mathcal{N} := \{x : S(x) = 1_{\mathbb{A}}\}$ is linear if, for all large v , one of $\pm v = T(u)$ is soluble.

Theorem (Maximal linear subspace characterization). *With assumptions as before,*

$$\mathcal{H} := \{x \in \mathbb{A} : (\forall t \in \mathbb{R}) N(tx) = tN(x)\} = \{u \in \mathbb{A} : S'(0)u = 0\} \subseteq \mathcal{N}.$$

So if \mathcal{N} is linear, then $\mathcal{N} = \mathcal{H}$ and N is linear on \mathcal{N} . For \mathcal{M} complementary to \mathcal{N} with $\pi_{\mathcal{M}}, \pi_{\mathcal{N}}$ corresponding projections:

$$N(x) = N(\pi_{\mathcal{N}}(x)) + N(\pi_{\mathcal{M}}(x)). \quad (+)$$

Key Tools: $\mathbb{G}_S(\mathbb{A})$ a Popa group under $x \circ_S y := x + S(x)y$ and the symbolic calculus. See [1].

References

- [1] Bingham, Nicholas Hugh, and Adam J. Ostaszewski. "The Gołab-Schinzel and Goldie functional equations in Banach algebras." *arXiv* (2021): arxiv.org/pdf/2105.07794.pdf.
- [2] Brillouët, Nicole, and Jean G. Dhombres. "Équations fonctionnelles et recherche de sous-groupes." *Aequationes Math.* 31, no. 2-3 (1986): 253-293.

Zdzisław Otachel

University of Life Sciences in Lublin, Poland

New advances on some inequalities in inner product spaces

Probably, the most known inequality related to inner product spaces is Cauchy-Schwarz inequality which states that the module of inner product of two vectors is not greater than product of norms of that vectors. The inequality is closely related to other, more sophisticated, inequalities of that type as Bessel's, Grüss', Gram's or Hadamard's inequalities.

The theory of such inequalities plays an important role in modern mathematics together with numerous applications for Nonlinear Analysis, Approximation and Optimization Theory, especially Engineering Optimization, Numerical Analysis, Probability Theory, Statistics, Theoretical Physics and other fields. The Schwarz and Grüss inequalities have been frequently used for obtaining bounds or estimating the errors in various approximation formulas occurring in the above domains. Thus, any new advancement will have a number of important consequences in the mathematical fields where inequalities are basic tools. New versions of classic inequalities also stimulates further developments of the more general theory of functional equations and inequalities.

The presentation basing on papers [1] - [5] is mainly devoted to reverses of Cauchy-Schwarz's inequality and consequences for other related inequalities. Cauchy-Schwarz inequality is valid for arbitrary vectors of the space, whereas its reverses hold true under certain restrictions and usually define upper bounds for the product of (squared) norms of two vectors reduced by (squared) module of inner product of these vectors. Applications to reverse Bessel's inequality, the reverse triangle inequality for norms and Grüss type inequalities are given and refinements of the famous Hadamard's inequality about determinants are presented.

References

- [1] Otachel, Zdzisław. "Inequalities for angles between subspaces with applications to Cauchy-Schwarz inequality in inner product spaces." *Math. Inequal. Appl.* 23, no. 2 (2020): 487-495.
- [2] Otachel, Zdzisław. "On Pólya-Szegő reverse of Schwarz's inequality." *Math. Nachr.* 293, no. 7 (2020): 1390-1404.
- [3] Otachel, Zdzisław. "Equality conditions of reverse Schwarz and Grüss inequalities in inner productspaces." *J. Math. Inequal.* 13, no. 2 (2019): 565-583.
- [4] Otachel, Zdzisław. "On reverse Schwarz, Bessel, Grüss and Hadamard inequalities in inner product spaces." *Linear Multilinear Algebra* 68, no. 4 (2020): 805-827.
- [5] Otachel, Zdzisław. "Reverse Schwarz inequalities and some consequences in inner product spaces." *J. Math. Anal. Appl.* 458, no. 2 (2018): 1409-1426.

Zsolt Páles 

University of Debrecen, Hungary

Limit theorems for deviation means of independent and identically
distributed random variables
(joint work with Mátyas Barczy)

Motivated by the Kolmogorov Strong Law of Large Numbers and the Lindberg–Lévy Central Limit Theorem, furthermore by some recent results of de Carvalho [1] concerning quasiarithmetic means and Barczy–Burai [2] concerning Bajraktarević means, replacing the arithmetic mean of a sequence of independent and identically distributed random variables by a deviation mean (cf. [3]), we obtain generalizations of the above mentioned limit theorems.

References

- [1] de Carvalho, Miguel. "Mean, what do you mean?" *Amer. Statist.* 70, no. 3 (2016): 270-274.
- [2] Barczy, Mátyás, and Pál Burai. "Limit theorems for Bajraktarević and Cauchy quotient means of independent identically distributed random variables." *Aequat. Math.* (2021). <https://doi.org/10.1007/s00010-021-00813-x>
- [3] Daróczy, Zoltán. "Über eine Klasse von Mittelwerten." *Publ. Math. Debrecen* 19 (1972): 211-217.

Paweł Pasteczka

Pedagogical University of Krakow, Poland

On the Riemannian approach to integral means

Adapting the notion of Riemann integral, for every weighted mean we define the corresponding lower and upper integral means. We show that such approach preserves most of properties of means, for example monotonicity and convexity/concavity.

References

- [1] Pasteczka, Paweł. "On the integral approach to means and their Hardy property." *arXiv* (2021): <https://arxiv.org/pdf/2003.10953.pdf>.

Kallol Paul 

Jadavpur University, Kolkata, India

k -smoothness of operators on infinite-dimensional spaces

In this talk we characterize k -smoothness of bounded linear operators defined between infinite-dimensional Hilbert spaces. The characterization of k -smoothness of bounded linear operators on arbitrary Banach space is still elusive. We try to address the problem in the setting of both finite and infinite-dimensional Banach spaces. This talk is based on the articles [1], [2].

References

- [1] Mal, Arpita, and Kallol Paul. "Characterization of k -smooth operators between Banach spaces." *Linear Algebra Appl.* 586 (2020): 296-307.
- [2] Mal, Arpita, Subhrajit Dey, and Kallol Paul. "Characterization of k -smoothness of operators defined between infinite-dimensional spaces," *Linear Multilinear Algebra* (2021). <https://doi.org/10.1080/03081087.2020.1844130>.

Adrian Petruşel

Babeş-Bolyai University Cluj-Napoca, Romania

Existence and Stability Results in Fixed Point Theory

(joint work with Gabriela Petruşel)

The purpose of this talk is to present some new results in metric fixed point theory for single-valued and multi-valued operators. The case of graphical (orbital) contractions will be considered. We will discuss existence of the fixed point, as well as data dependence, well-posedness, Ostrowski stability property and Ulam-Hyers stability of the fixed point problem. In the case of the stability properties associated to the fixed point problem, the approach will put in the light the role of the retraction-displacement condition.

References

- [1] Feng, Yu Qiang, and San Yang Liu. "Fixed point theorems for multi-valued contractive mappings and multi-valued Caristi type mappings." *J. Math. Anal. Appl.* 317, no. 1 (2006): 103-112.
- [2] Minak, Gülhan, Özlem Acar, and İshak Altun. "Multivalued pseudo-Picard operators and fixed point results." *J. Funct. Spaces Appl.* (2013): Article ID 827458.
- [3] Petruşel, Adrian, Gabriela Petruşel, and Jen-Chih Yao. "Multivalued graph contraction principle." *Optimization* 69, no. 7-8 (2020): 1541-1556.
- [4] Petruşel, Adrian, and Ioan Aurel Rus. "Graphic contraction principle and applications." In: *Mathematical Analysis and Applications* Rassias T., Pardalos P. (eds), 411-432. Cham: Springer, 2019
- [5] Petruşel, Adrian, and Gabriela Petruşel. "Some variants of the contraction principle for multi-valued operators, generalizations and applications." *J. Nonlinear Convex Anal.* 20, no. 10 (2019): 2187-2203.
- [6] Rheinboldt, Werner. "A unified convergence theory for a class of iterative processes." *SIAM J. Numer. Anal.* 5 (1968): 42-63.
- [7] Subrahmanyam, Papagudi Venkatachalam. "Remarks on some fixed-point theorems related to Banach's contraction principle." *J. Mathematical and Physical Sci.* 8 (1974): 445-457; errata 9 (1975): 195.

Dorian Popa

Technical University of Cluj-Napoca, Romania

Best constant in Ulam Stability

In this talk I present results on the best Ulam constant for some functional equations and some linear operators.

Teresa Rajba
University of Bielsko-Biala, Poland

On the Raşa inequality for higher-order convex functions

This talk is based on the article [2]. We study the following $(q - 1)$ th convex ordering relation for q th convolution power of the difference of probability distributions μ and ν

$$(\nu - \mu)^{*q} \geq_{(q-1)cx} 0, \quad q \geq 2,$$

and we obtain the theorem providing a useful sufficient condition for its verification. We apply this theorem for various families of probability distributions and we obtain several inequalities related to the classical interpolation operators. In particular, taking binomial distributions, we obtain a new, very short proof of the inequality given recently by Abel and Leviatan [1].

Theorem 1 ([1]). *Let $q, n \in \mathbb{N}$, $q \geq 2$ and $x, y \in [0, 1]$. If $f \in \mathcal{C}([0, 1])$ is a $(q - 1)$ -convex function, then*

$$\begin{aligned} \operatorname{sgn}(x - y)^q \sum_{\nu_1, \dots, \nu_q=0}^n \sum_{j=0}^q (-1)^{q-j} \binom{q}{j} \left(\prod_{i=1}^j p_{n, \nu_i}(x) \right) \left(\prod_{i=j+1}^q p_{n, \nu_i}(y) \right) \\ \times f\left(\frac{\nu_1 + \dots + \nu_q}{qn}\right) \geq 0, \quad (1) \end{aligned}$$

where

$$p_{n,i}(x) = \binom{n}{i} x^i (1 - x)^{n-i}, \quad 0 \leq i \leq n.$$

Inequality (1) is a generalization of inequality stated by Ioan Raşa as an open problem in [4] and proved in [3].

Theorem 2 ([3]). *Let $n \in \mathbb{N}$ and $x, y \in [0, 1]$. Then*

$$\sum_{i=0}^n \sum_{j=0}^n (p_{n,i}(x)p_{n,j}(x) + p_{n,i}(y)p_{n,j}(y) - 2p_{n,i}(x)p_{n,j}(y)) f\left(\frac{i+j}{2n}\right) \geq 0$$

for all convex functions $f \in \mathcal{C}([0, 1])$.

References

- [1] Abel, Ulrich, and Dany Leviatan. "An extension of Raşa's conjecture to q -monotone functions." *Results Math.* 75, no. 4 (2020): Paper No. 180.
- [2] Komisarski, Andrzej, and Teresa Rajba. "On the Raşa inequality for higher order convex functions." *Results Math.* 76, no. 2 (2021): Paper No. 103.
- [3] Mrowiec, Jacek, Teresa Rajba, and Szymon Wąsowicz. "A solution to the problem of Raşa connected with Bernstein polynomials." *J. Math. Anal. Appl.* 446, no. 1 (2017): 864-878.
- [4] Raşa, Ioan. "Problem, p. 164." In: *Report of Meeting "Conference on Ulam's Type Stability", Rytro, Poland, June 2-6, 2014*. Ann. Univ. Paedagog. Crac. Stud. Math. 13 (2014): 139-169.

Ioan Raşa 

Technical University of Cluj-Napoca, Romania

A functional equation related to Appell polynomials and Heun functions (joint work with Ana Maria Acu)

We are concerned with functional equations of the form

$$\omega(x)y'_n(x) = n\omega'(x)(y_n(x) - y_{n-1}(x)), \quad (1)$$

where ω is a given function on an interval I . First, using the theory of Appell polynomials we determine all the solutions of (1). Then we present equations of type (1) for which the solutions are log-convex. Some polynomial Heun functions are log-convex solutions of such equations. We investigate the properties of the involved Appell polynomials and apply the results to the associated Jakimovski-Leviatan operators. Our results extend and generalize results from [1]-[4].

References

- [1] Acu, Ana Maria, Ioan Cristian Buscu, and Ioan Raşa. "A sequence of Appell polynomials and the associated Jakimovski-Leviatan operators." *Anal. Math. Phys.* 11, no. 2 (2021): Paper No. 88.
- [2] Alzer, Horst. "Remarks on a convexity theorem of Raşa." *Results Math.* 75, no. 1 (2020): Paper No. 29.
- [3] Nikolov, Geno. "Inequalities for ultraspherical polynomials. Proof of a conjecture of I. Raşa." *J. Math. Anal. Appl.* 418, no. 2 (2014): 852-860.
- [4] Raşa, Ioan. "Convexity properties of some entropies (II)." *Results Math.* 74, no. 4 (2019): Paper No. 154.

Lenka Rucká

Silesian university in Opava, Czech Republic

Some results on distributionally chaotic points (joint work with Francisco Balibrea)

The distributionally chaotic point (DC point for short) is introduced in [1] as a point whose arbitrarily small neighbourhood contains an uncountable distributionally chaotic set, which is bounded by a special envelope. In this talk we extend the result from [1] and we show that for continuous interval maps, positive topological entropy implies existence of uncountably many DC points. Also we show that this result cannot be extended to higher class of maps, particularly to continuous triangular maps of the square.

References

- [1] Pawlak, Ryszard Jerzy, and Anna Loranty. "On the local aspects of distributional chaos." *Chaos* 29, no. 1 (2019): 013104.

Maciej Sablik
University of Silesia, Katowice, Poland

On some new functional equations characterizing polynomial functions
(joint work with Chisom Prince Okeke)

We are considering the following class of equations

$$F(x+y) - F(x) - F(y) = \sum_{i=1}^m (a_i x + b_i y) f(\alpha_i x + \beta_i y), \quad (1)$$

where $F, f: \mathbb{R} \rightarrow \mathbb{R}$ are unknown functions and for each $i \in \{1, \dots, m\}$ the numbers $a_i, b_i \in \mathbb{R}$ and $\alpha_i, \beta_i \in \mathbb{Q}$ are given.

Equation (1) is a far going extension of the equation considered by Włodzimierz Fechner and Eszter Gselmann, namely

$$F(x+y) - F(x) - F(y) = xf(y) + yf(x), \quad (2)$$

which was studied by the authors in [1]. Together with Timothy Nadjohi and Tomasz Szostok we studied in [2] the following extension of (2)

$$\sum_{i=1}^n \gamma_i F(\alpha_i x + \beta_i y) = xf(y) + yf(x). \quad (3)$$

The results have been published in [2].

We present a method of solving equations of type (1), based on a Lemma contained in [2].

This is the theoretical part of the talk which has been divided into two parts. The second will be presented by Chisom Prince Okeke.

References

- [1] Fechner, Włodzimierz, and Eszter Gselmann. "General and alien solutions of a functional equation and of a functional inequality." *Publ. Math. Debrecen* 80, no. 1-2 (2012): 143-154.
- [2] Nadjohi, Timothy, Chisom Prince Okeke, Maciej Sablik, and Tomasz Szostok. "On a new class of functional equations satisfied by polynomial functions." *Aequat. Math.* (2021). <https://doi.org/10.1007/s00010-021-00781-2>

Debmalya Sain 

Indian Institute of Science, Bangalore, India

Best approximations from the perspective of orthogonality

We plan to discuss an application of Birkhoff-James orthogonality in studying the best approximation problem in Banach spaces. This allows us to obtain some interesting distance formulae for both compact operators and vectors. The advantage of our method over the classical duality principle, especially from a computational point of view, will also be discussed. This talk is based on my recent article [1]

References

- [1] Sain, Debmalaya. "On best approximations to compact operators." *Proc. Amer. Math. Soc.* 149, no. 10 (2021): 4273-4286.

Adolf Schleiermacher

no affiliation

On even orthogonally additive mappings (joint work with Justyna Sikorska)

In 1990 György Szabó proved: *If an orthogonality space (X, \perp) in the sense of Rätz [3] and of dimension at least 3 over the field of real numbers admits a nontrivial even orthogonally additive mapping $E: X \rightarrow Y$ into some Abelian group Y , then X is equivalent to an inner product space (cf. [4]).* Using methods from projective geometry (cf. [1], [2]) we prove a stronger theorem consisting of several equivalent assertions about (X, \perp) and valid for arbitrary ordered groundfields K . Note that Dilian Yang has proved in [5] an analogue of Szabó's theorem for the case of dimension two. However, this case requires different methods.

References

- [1] Baer, Reinhold. *Linear Algebra and Projective Geometry*. New York: Academic Press, 1952.
- [2] Lenz, Hanfried. "Über die Einführung einer absoluten Polarität in die projektive und affine Geometrie des Raumes" *Math. Ann.* 128 (1954): 363-372.
- [3] Rätz, Jürg. "On orthogonally additive mappings." *Aequationes Math.* 28, no. 1-2 (1985): 35-49.
- [4] Szabó, György. "On orthogonality spaces admitting nontrivial even orthogonally additive mappings." *Acta Math. Hungar.* 56, no. 1-2 (1990): 177-187.
- [5] Yang, Dilian. "Orthogonality spaces and orthogonally additive mappings." *Acta Math. Hungar.* 113, no. 4 (2006): 269-280.

A class of functional equations associated with almost periodic functions

During this talk we shortly describe how to get a class of functional equations involving a countable set of terms, summed by the well known Bochner-Fejér summation procedure, which are closely associated with the set of Bohr's almost periodic functions. First, we discuss the case of a finite number of terms in connection with exponential polynomials, including the initial motivation with the partial sums of the Riemann zeta function. Secondly, in conjunction with the study of some particular properties and solutions, we progressively extend the expression of our functional equations to the case of a countable set of terms, and we finally show that the zeros of a prefixed almost periodic function determine analytic solutions of such a functional equation associated with it. This talk is based on the articles [1], [2], [3] and [4].

References

- [1] Mora Martínez, Gaspar, and Juan Matías Sepulcre. "The zeros of Riemann zeta partial sums yield solutions to $f(x) + f(2x) + \dots + f(nx) = 0$." *Mediterr. J. Math.* 10, no. 3 (2013): 1221-1233.
- [2] Sepulcre, Juan Matías. "On some solutions of a functional equation related to the partial sums of the Riemann zeta function." *Bull. Korean Math. Soc.* 51, no. 1 (2014): 29-41.
- [3] Sepulcre, Juan Matías, and Tomás Vidal. "On the analytic solutions of the functional equations $w_1 f(a_1 z) + w_2 f(a_2 z) + \dots + w_n f(a_n z) = 0$." *Mediterr. J. Math.* 12, no. 3 (2015): 667-678.
- [4] Sepulcre, Juan Matías, and Tomás Vidal. "A class of functional equations associated with almost periodic functions." *Aequationes Math.* 95, no. 1 (2021): 91-105.

Periodic points of asymmetric Bernoulli shifts

In 1964, Sharkovskii [4] firstly introduced a special ordering on the set of positive integers. This ordering implies that if $p \triangleleft q$ and a self-map of a closed bounded interval has a point of period p , then it has a point of period q . The least number with respect to this ordering is 3. Thus, if a map has a point of period 3, then it has points of any periods. In 1975, the latter result was rediscovered by Li and Yorke [1]. Then numerous papers are devoted to the study of interval maps (see, e.g. [2], [3], [5] and references therein)

Consider asymmetric Bernoulli shift $F: [0, 1] \rightarrow [0, 1]$ with a parameter $0 < a < 1$, defined by

$$F(x) := \begin{cases} \frac{x}{a}, & 0 \leq x \leq a, \\ \frac{x-a}{1-a}, & a < x \leq 1. \end{cases}$$

Given a positive integer n , one interesting question is how to find all n -periodic points of F . The other is how many n -periodic points of F .

References

- [1] Li, Tien-Yien, and James Alan Yorke. "Period three implies chaos." *Amer. Math. Monthly* 82, no. 10 (1975): 985-992.
- [2] Milnor, John, and William Thurston. "On Iterated Maps of the Interval." In: *Dynamical Systems: Proceedings 1986-1987* Vol. 1342 of *Lecture Notes in Mathematics*. Berlin-Heidelberg-New York: Springer-Verlag, 1988: 465-563.
- [3] Glendinning, Paul. "Topological conjugation of Lorenz maps by β -transformations." *Math. Proc. Cambridge Philos. Soc.* 107, no. 2 (1990): 401-413.
- [4] Sharkovsky, Oleksandr Mykolayovych. "Co-existence of cycles of a continuous mapping of the line into itself" *Ukrain. Mat. Ž.* 16 (1964): 61-71.
- [5] Bassein, Susan. "The dynamics of a family of one-dimensional maps." *Amer. Math. Monthly* 105, no. 2 (1998): 118-130.

Justyna Sikorska

University of Silesia, Katowice, Poland

On a generalized n -linear equation (joint work with Anna Bahyrycz)

General linear functional equations have been studied for years. During the talk we shall discuss their counterpart for multivariable functions.

Let X, Y be linear spaces over a field \mathbb{K} and $f: X^n \rightarrow Y$. For some fixed $a_{ji} \in \mathbb{K} \setminus \{0\}$, $C_{i_1 \dots i_n} \in \mathbb{K}$ for all $j \in \{1, \dots, n\}$, $i, i_k \in \{1, 2\}$, $k \in \{1, \dots, n\}$, we consider the following equation

$$f(a_{11}x_{11} + a_{12}x_{12}, \dots, a_{n1}x_{n1} + a_{n2}x_{n2}) = \sum_{1 \leq i_1, \dots, i_n \leq 2} C_{i_1 \dots i_n} f(x_{1i_1}, \dots, x_{ni_n}), \quad (*)$$

for all $x_{ji_j} \in X$, $j \in \{1, \dots, n\}$, $i_j \in \{1, 2\}$.

We determine the general solution of $(*)$ and as special cases we obtain some results for multi-Cauchy, multi-Jensen and multi-Cauchy-Jensen equations.

Global bounds for the generalized Jensen functional with applications (joint work with Vesna Todorčević)

Recall that the Jensen functional $J_n(\mathbf{p}, \mathbf{x}; h)$ is defined on an interval $I \subseteq \mathbb{R}$ by

$$J_n(\mathbf{p}, \mathbf{x}; h) := \sum_{i=1}^n p_i h(x_i) - h\left(\sum_{i=1}^n p_i x_i\right),$$

where $h: I \rightarrow \mathbb{R}$, $\mathbf{x} = (x_1, x_2, \dots, x_n) \in I^n$ and $\mathbf{p} = \{p_i\}_{i=1}^n$ is a positive weight sequence.

If h is a convex function on I , then the inequality $0 \leq J_n(\mathbf{p}, \mathbf{x}; h)$ holds for each $\mathbf{x} \in I^n$ and any positive weight sequence \mathbf{p} .

Supposing that $\mathbf{x} \in [a, b]^n \subset I^n$, our aim in this paper is to find some *global bounds* for the generalized Jensen functional

$$J_n(\mathbf{p}, \mathbf{x}; g, h) := g\left(\sum_{i=1}^n p_i h(x_i)\right) - g\left(h\left(\sum_{i=1}^n p_i x_i\right)\right),$$

i.e. the bounds which does not depend on \mathbf{p} or \mathbf{x} but only on a, b and functions g and h .

In this sense, a typical result is given by the following

Theorem. *Let $h: I \rightarrow J$ be convex and $g: J \rightarrow \mathbb{R}$ be an increasing function. Then*

$$0 \leq J_n(\mathbf{p}, \mathbf{x}; g, h) \leq \max_p [g(ph(a) + (1-p)h(b)) - g(h(pa + (1-p)b))].$$

We give some applications regarding $\mathcal{A} - \mathcal{G} - \mathcal{H}$ inequalities, where

$$\mathcal{A}(\mathbf{p}, \mathbf{x}) := \sum p_i x_i; \quad \mathcal{G}(\mathbf{p}, \mathbf{x}) := \prod x_i^{p_i}; \quad \mathcal{H}(\mathbf{p}, \mathbf{x}) := \left(\sum p_i / x_i\right)^{-1},$$

are the generalized arithmetic, geometric and harmonic means, respectively.

For example, let $\mathbf{x} \in [a, b]^n$, $0 < a < b$. Then $0 \leq \mathcal{A}(\mathbf{p}, \mathbf{x}) - \mathcal{H}(\mathbf{p}, \mathbf{x}) \leq 2(A(a, b) - G(a, b))$;

$$1 \leq \frac{\mathcal{A}(\mathbf{p}, \mathbf{x})}{\mathcal{H}(\mathbf{p}, \mathbf{x})} \leq \left(\frac{A(a, b)}{G(a, b)}\right)^2; \quad 1 \leq \frac{\mathcal{G}(\mathbf{p}, \mathbf{x})}{\mathcal{H}(\mathbf{p}, \mathbf{x})} \leq \frac{I(a, b)L(a, b)}{G^2(a, b)};$$

$$0 \leq \mathcal{A}(\mathbf{p}, \mathbf{x}) - \mathcal{G}(\mathbf{p}, \mathbf{x}) \leq A(a, b) - L(a, b) + L(a, b) \log \frac{L(a, b)}{G(a, b)}; \quad 1 \leq \frac{\mathcal{A}(\mathbf{p}, \mathbf{x})}{\mathcal{G}(\mathbf{p}, \mathbf{x})} \leq \frac{I(a, b)L(a, b)}{G^2(a, b)},$$

where A, G, H, L, I stands for the arithmetic, geometric, harmonic, logarithmic and intrinsic means of positive numbers a and b , respectively. All bounds above are best possible.

Open problem. Find the best possible global converse for $\mathcal{G}(\mathbf{p}, \mathbf{x}) - \mathcal{H}(\mathbf{p}, \mathbf{x})$.

References

- [1] Simić, Slavko. "Sharp global bounds for Jensen's inequality" *Rocky Mountain J. Math.* 41, no. 6 (2011): 2021-2031.

László Székelyhidi
University of Debrecen, Hungary

Functional equations, moment functions and Bell polynomials
(joint work with Żywilla Fechner and Eszter Novák-Gselmann)

During this talk we present how to describe moment function sequences of higher rank on commutative groups and on polynomial hypergroups. This talk is based on the articles [2] and [3]. The description depends on certain functional equation systems which generalize some binomial-type functional equation systems considered and solved in [1]. For the general description we use Bell polynomials.

References

- [1] Aczél, János. "Functions of binomial type mapping groupoids into rings." *Math. Z.* 154, no. 2 (1977): 115-124.
- [2] Fechner, Żywilla, Eszter Gselmann, and László Székelyhidi. "Moment functions on groups." *Results Math.* 76 (2021): Article No. 171.
- [3] Fechner, Żywilla, Eszter Gselmann, and László Székelyhidi. "Moment functions and exponential monomials on commutative hypergroups." *Aequat. Math.* (2021). <https://doi.org/10.1007/s00010-021-00827-5>

Patricia Szokol

University of Debrecen, Hungary

Hermite - Hadamard type inequality for certain Schur convex functions (joint work with Pál Burai and Judit Makó)

In the presentation we investigate symmetric, continuous n -variable functions that satisfy a property, which can be considered as a Hermite-Hadamard type inequality. In the main result we prove that such functions are necessarily Jensen-convex. It turns out, that the key tool of the proof is a Korovkin-type approximation theorem, which will be presented in the talk, as well. Finally, we present some applications of our main theorem.

References

- [1] Bessenyei, Mihály, and Zsolt Páles. "Characterization of convexity via Hadamard's inequality." *Math. Inequal. Appl.* 9, no. 1 (2006): 53-62.
- [2] Burai, Pál, and Judit Makó. "On certain Schur-convex functions." *Publ. Math. Debrecen* 89, no. 3 (2016): 307-319.
- [3] Burai, Pál, Judit Makó, and Patrícia Szokol. "Hermite-Hadamard type inequality for certain Schur convex functions." *Math. Inequal. Appl.* 23, no. 4 (2020): 1441-1458.
- [4] Hadamard, Jacques. "Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann." *J. Math. Pures Appl.* 58 (1893): 171-215.
- [5] Gilányi, Attila, Kazimierz Nikodem, and Zsolt Páles. "Bernstein-Doetsch type results for quasicontinuous functions." *Math. Inequal. Appl.* 7, no. 2 (2004): 169-175.
- [6] Korovkin, Pavel Petrovich. "On convergence of linear positive operators in the space of continuous functions." *Doklady Akad. Nauk SSSR (N.S.)* 90 (1953): 961-964.
- [7] Makó, Judit, and Attila Házy. "On approximate Hermite-Hadamard type inequalities." *J. Convex Anal.* 24, no. 2 (2017): 349-363.

Tomasz Szostok

University of Silesia, Katowice, Poland

On some class of means

We consider means connected with quadrature rules of numerical integration. Then the functions which generate weighted arithmetic mean in this way are characterized and some inequalities between different means of this type are proved.

Lan Nhi To

University of Debrecen, Hungary

Computer assisted investigation of the alienness of linear functional equations (joint work with Attila Gilányi)

The concept of the alienness and the strong alienness of functional equations was introduced by J. Dhombres in [3]. The properties were investigated by many authors during the last years (cf., e.g. [4], [5], [7]).

In this talk, we present a computer program, developed in the computer algebra system Maple, which investigates the alienness and the strong alienness of linear functional equations of the type

$$\sum_{i=0}^{n+1} f_i(p_i x + q_i y) = 0 \quad (x, y \in X),$$

where n is a positive integer, p_0, \dots, p_{n+1} and q_0, \dots, q_{n+1} are rational numbers, X, Y are linear spaces and $f_0, \dots, f_{n+1}: X \rightarrow Y$ are unknown functions.

Our application is based on a computer program developed for the solution of linear functional equations of the type above, presented in [6] (cf. also [1] and [2]).

References

- [1] Borus, Gergő Gyula, and Attila Gilányi. "Solving systems of linear functional equations with computer." *2013 IEEE 4th International Conference on Cognitive Infocommunications (CogInfoCom)*, 559-562. Budapest, 2013.
- [2] Borus, Gergő Gyula, and Attila Gilányi. "Computer assisted solution of systems of two variable linear functional equations." *Aequationes Math.* 94, no. 4 (2020): 723-736.
- [3] Dhombres, Jean G. "Relations de dépendance entre les équations fonctionnelles de Cauchy." *Aequationes Math.* 35, no. 2-3 (1988): 186-212.
- [4] Ger, Roman. "On alienation of two functional equations of quadratic type", *Aequat. Math.* (2021). <https://doi.org/10.1007/s00010-021-00809-7>
- [5] Ger, Roman, and Maciej Sablik. "Alien functional equations: a selective survey of results." In: *Developments in Functional Equations and Related Topics* Vol. 124 of *Springer Optimization and Its Applications*. Cham: Springer, 2017: 107-147.
- [6] Gilányi, Atilla. *Charakterisierung von monomialen Funktionen und Lösung von Funktionalgleichungen mit Computern*. Karlsruhe: Universität Karlsruhe, 1995.
- [7] Szostok, Tomasz. "Alienation of two general linear functional equations." *Aequationes Math.* 94, no. 2 (2020): 287-301.

Radu Truşcă

Babeş-Bolyai University, Cluj-Napoca, Romania

Some local fixed point results and applications for generalized contractions

The purpose of this work is to present some local fixed point results, under weaker assumptions, for three types of generalized contractions: Ćirić-Reich-Rus, Chatterjea and Berinde. As applications we will present open mapping principles and continuation results for the three aforementioned classes of contractions. For example, let us consider the class of Ćirić-Reich-Rus generalized contractions.

Definition . Let (X, d) be a metric space. We say that an operator $f: X \rightarrow X$ is a *Ćirić-Reich-Rus contraction* if there exist $\alpha, \beta \in \mathbb{R}^+$ with $\alpha + 2\beta < 1$ such that we have

$$d(f(x), f(y)) \leq \alpha d(x, y) + \beta [d(x, f(x)) + d(y, f(y))], \text{ for all } x, y \in X. \quad (1)$$

The local fixed point theorem related to Ćirić-Reich-Rus contractions is the following one.

Theorem 1. Let (X, d) be a complete metric space, $x_0 \in X$, a positive number r and let $f: B(x_0; r) \rightarrow X$ be a Ćirić-Reich-Rus type contraction. If

$$d(x_0, f(x_0)) < \frac{1 - \alpha - 2\beta}{1 - \beta} r, \quad (2)$$

then the sequence $(f^n(x_0))_{n \in \mathbb{N}}$ converges to a point x^* which is a fixed point for the Ćirić-Reich-Rus contraction f . Moreover, the fixed point is unique.

As applications of our result, we will give open mapping principles and continuation results.

References

- [1] Berinde, Vasile, and Mădălina Păcurar. "Iterative approximation of fixed points of single-valued almost contractions". In: *Fixed Point Theory and Graph Theory*. Amsterdam: Elsevier–Academic Press: 29-97, 2016.
- [2] Chatterjea, Santi Kumar. "Fixed-point theorems." *C. R. Acad. Bulgare Sci.* 25 (1972): 727-730.
- [3] Ćirić, Ljubomir B.. "Generalized contractions and fixed-point theorems." *Publ. Inst. Math. (Beograd) (N.S.)* 12(26) (1971): 19-26.
- [4] Granas, Andrzej and James Dugundji. *Fixed Point Theory*. Springer Monographs in Mathematics. Berlin: Springer, 2003.
- [5] Lazăr, Tania Angelica, and Naseer Shahzad. "Fixed points for non-self operators and domain invariance theorems." *Nonlinear Anal.* 70, no. 1 (2009): 117-125.
- [6] Reich, Simeon. "Fixed points of contractive functions." *Unione Mat. Ital.* 4 (1972): 26-42.
- [7] Truşcă, Radu. "Some local fixed point theorems and applications to open mapping principles and continuation results." *Arab. J. Math* (2021): to appear. <https://doi.org/10.1007/s40065-021-00331-3>

On preserving the relaxed polygonal inequality

During the presentation we investigate the families of functions of the form $f: [0, +\infty) \rightarrow [0, +\infty)$ which preserve the relaxed polygonal inequality, introduced by Fagin et al. in [4]. Formally speaking, the definition of this concept is as follows:

Definition. Relaxed polygonal inequality

Let (X, d) be a semimetric space. If there exists $\gamma \geq 1$ such that

$$\forall_{n \in \mathbb{N}} \quad \forall_{x_1, \dots, x_n \in X} \quad d(x_1, x_n) \leq \gamma \cdot \sum_{k=1}^{n-1} d(x_k, x_{k+1}), \quad (1)$$

then we say that d satisfies the *relaxed polygonal inequality* (*rpi* in short) with the relaxation constant γ .

We investigate the families of functions which can be composed with any semimetric satisfying rpi to obtain another semimetric with this property (such mappings will be called (P) -preserving). In general, the concept of property-preserving functions was introduced as early as in 1947 by Sreenivasan [5], but most profound results were formulated by Slovakian mathematicians – Borsík and Doboš (e.g. [1], [2], [3]).

We would like to discuss two theorems which give equivalent conditions for f to be (P) -preserving.

Firstly, we introduce the easily-verifiable characterization for increasing (P) -preserving functions. Then we introduce some special finite tuples of positive real numbers which are meant to resemble distances between points in some semimetric space satisfying rpi. These so-called γ -polygons will be then used in our second characterization of (P) -preserving mappings. In particular, we will prove that every such mapping has to satisfy $f^{-1}[\{0\}] = \{0\}$ and for any $K_1 \geq 1$, there exists K_2 such that f maps every K_1 -relaxed polygon to K_2 -relaxed polygon.

References

- [1] Borsík, Ján, and Jozef Doboš. "On metric preserving functions." *Real Anal. Exchange* 13, no. 1 (1987/88): 285-293.
- [2] Doboš, Jozef. *Metric preserving functions..* Kosice: Stroffek, 1998. <http://web.science.upjs.sk/jozefdobos/wp-content/uploads/2012/03/mpf1.pdf>
- [3] Doboš, Jozef. "On modifications of the Euclidean metric on reals." *Tatra Mt. Math. Publ.* 8 (1996): 51-54.
- [4] Fagin, Ronald, Ravi Kumar, and Dandapani Sivakumar. "Comparing top k lists." *SIAM J. Discrete Math.* 17, no. 1 (2003): 134-160.
- [5] Sreenivasan, T. K. "Some properties of distance functions." *J. Indian Math. Soc. (N.S.)* 11 (1947): 38-43.

Szymon Wąsowicz 
University of Bielsko-Biala, Poland

New adaptive method of the approximate integration

The classical Simpson's composite rule induces an adaptive method of the approximate integration, which stops an algorithm whenever some control expression containing the fourth derivative of an integrand is less from a desired precision of computation. Sometimes it is difficult to estimate this derivative. Then finding an algorithm which does not require any derivatives seems to be useful. Clenshaw and Curtis described in 1960 (without any proof) and Rowland and Varol examined in 1972 a method based precisely on the Simpson's rule. This is a starting point of our considerations. In the talk we introduce another method, which involves some convex combination of the Simpson's rule and the Chebyshev quadrature. The objects of our research are so-called convex functions of order three. We present the results of the numerical experiments which convince us that our adaptive method requires considerably less subpartitions of the interval of integration than the classical methods.

Bettina Wilkens
University of Namibia

Algebraic methods in spectral synthesis on discrete abelian groups

During this talk we present some applications of classical results of commutative algebra to established results and open problems in the area of spectral synthesis on discrete abelian groups. We can say more. Let G be an Abelian group and let V be a variety in \mathcal{CG} . *Spectral synthesis holds on V* if every subvariety is the closure of its finite-dimensional subvarieties. We can show something more:

Theorem. *Let V be a variety on the Abelian group G . Let $R = \mathbb{C}G/V^\perp$. For a maximal ideal M of R , let $W_M = \bigcup_{n \in \mathbb{N}} (M^n)^\perp$. If spectral synthesis holds on V , then:*

1. *The variety V is spanned by the varieties W_M , M running over the maximal ideals of R .*
2. *For each M , W_M has a subvariety U_M spanned by finite-dimensional subvarieties annihilated by a power of M .*
3. *For each M , W_M is the closure of U together with subvarieties of dimension bounded by some integer N depending only on M .*

Let U be any R -module. We say that U is *finitary* if, for every $x \in R$ and $u \in U$, there is $0 \neq f \in \mathbb{C}[x]$ such that $uf = 0$.

Lemma. *If R has a faithful finitary module, then local spectral synthesis holds on V .*

Local spectral synthesis (see [2]) is the property of being the closure of the linear span of functions restricting to polynomials on any finitely generated subgroup. We finish by discussing potential converses of the above lemma.

References

- [1] Laczkovich, Miklós, and László Székelyhidi. "Spectral synthesis on discrete abelian groups." *Math. Proc. Cambridge Philos. Soc.* 143, no. 1 (2007): 103-120.
- [2] Laczkovich, Miklós. "Local spectral synthesis on Abelian groups." *Acta Math. Hungar.* 143, no. 2 (2014): 313-329.
- [3] Székelyhidi, László. "The failure of spectral synthesis on some types of discrete abelian groups." *J. Math. Anal. Appl.* 291, no. 2 (2004): 757-763.
- [4] Székelyhidi, László. "Annihilator methods in discrete spectral synthesis." *Acta Math. Hungar.* 143, no. 2 (2014): 351-366.
- [5] Székelyhidi, László. *Harmonic and Spectral Analysis*. Singapore, Hackensack, London: World Scientific Publishing Co. Pte. Ltd., 2014.
- [6] Székelyhidi, László, and Bettina Wilkens. "Spectral synthesis and residually finite-dimensional algebras." *J. Algebra Appl.* 16, no. 10 (2017): Article no. 1750200.

Paweł Wójcik

Pedagogical University of Krakow, Poland

Generalized orthogonality equation

(joint work with Tomasz Kania and Tomasz Kobos)

During this talk we show that every map between finite-dimensional normed spaces of the same dimension that respects fixed semi-inner products must be automatically a linear isometry. More precisely, the main results is as follows:

Theorem 1. *Let X and Y be normed spaces with fixed semi-inner products $[\cdot|\cdot]_X$, $[\cdot|\cdot]_Y$, respectively. Suppose that $f: X \rightarrow Y$ is a function such that*

$$[f(x)|f(y)]_Y = [x|y]_X, \quad x, y \in X.$$

If either

(a) X and Y have the same finite dimension, or

(b) X has a Schauder basis (e_i) and $(f(e_i))$ is a Schauder basis of Y ,

then f is a linear isometry.

Moreover, we construct a uniformly smooth renorming of the Hilbert space ℓ_2 and a continuous injection acting thereon that respects the semi-inner products, yet it is non-linear.

Sebastian Wójcik

University of Rzeszów, Poland

Quasi-arithmetic type means generated by the generalized Choquet integral

We introduce a class of quasi-arithmetic type means generated by the generalized Choquet integral. They are related to the notion of certainty equivalent under Cumulative Prospect Theory and play a significant role, e.g. in a utility-based insurance contracts pricing (cf. [2]). We establish the characterizations of the equality, positive homogeneity, and translativity for such means. In particular, we generalize some results in [1].

References

- [1] Chudziak, Jacek. "Certainty equivalent under cumulative prospect theory." *Internat. J. Uncertain. Fuzziness Knowledge-Based Systems* 27, no. 3 (2019): 415-428.
- [2] Kałuszką, Marek, and Michał Krzeszowiec. "Mean-value principle under cumulative prospect theory." *Astin Bull.* 42, no. 1 (2012): 103-122.

Amr Zakaria 

Ain Shams University, Cairo, Egypt

An invariance problem with six unknown functions
(joint work with Zsolt Páles)

Given two continuous functions $f, g: I \rightarrow \mathbb{R}$ such that g is nowhere zero on I and the ratio function f/g is strictly monotone on I , the *weighted two-variable Bajraktarević mean* $B_{f,g}: I^2 \times \mathbb{R}_+^2 \rightarrow I$ is defined by

$$B_{f,g}((x, y), (t, s)) := \left(\frac{f}{g}\right)^{-1} \left(\frac{tf(x) + sf(y)}{tg(x) + sg(y)} \right) \quad (x, y \in I; s, t \in \mathbb{R}_+).$$

The purpose of this talk is to solve the invariance of a Bajraktarević mean with respect to two weighted Bajraktarević means. In more details, we intend, under higher order regularity assumptions, to solve the functional equation

$$\frac{\varphi(B_{f,g}((x, y), (t, s))) + \varphi(B_{h,k}((x, y), (s, t)))}{\psi(B_{f,g}((x, y), (t, s))) + \psi(B_{h,k}((x, y), (s, t)))} = \frac{\varphi(x) + \varphi(y)}{\psi(x) + \psi(y)} \quad (x, y \in I),$$

or equivalently,

$$\left| \begin{array}{cc} \varphi(B_{f,g}((x, y), (t, s))) + \varphi(B_{h,k}((x, y), (s, t))) & \varphi(x) + \varphi(y) \\ \psi(B_{f,g}((x, y), (t, s))) + \psi(B_{h,k}((x, y), (s, t))) & \psi(x) + \psi(y) \end{array} \right| = 0 \quad (x, y \in I),$$

where $f, g, h, k, \varphi, \psi: I \rightarrow \mathbb{R}$ are unknown continuous functions such that g, k, ψ are nowhere zero on I , the ratio functions $f/g, h/k, \varphi/\psi$ are strictly monotone on I , and s, t are fixed different positive numbers. This invariance problem is more general than the functional equation investigated in [3].

References

- [1] Bajraktarević, Mahmud. "Sur une équation fonctionnelle aux valeurs moyennes." *Glasnik Mat.-Fiz. Astronom. Društvo Mat. Fiz. Hrvatske Ser. II* 13 (1958): 243-248.
- [2] Bajraktarević, Mahmud. "Über die Vergleichbarkeit der mit Gewichtsfunktionen gebildeten Mittelwerte." *Studia Sci. Math. Hungar.* 4 (1969): 3-8.
- [3] Páles, Zsolt, and Amr Zakaria. "On the invariance equation for two-variable weighted nonsymmetric Bajraktarević means." *Aequationes Math.* 93, no. 1 (2019): 37-57.

Marek Cezary Zdun
Pedagogical University of Krakow, Poland

On a linearization of Koopmans recursion and its application in economics

Let X be a metric space. Let \succeq be a binary relation on infinite product X^∞ and $U: X^\infty \rightarrow \mathbb{R}$ be a continuous surjection, such that $(x_0, x_1, x_2, \dots) \succeq (y_0, y_1, y_2, \dots) \Leftrightarrow U(x_0, x_1, x_2, \dots) \geq U(y_0, y_1, y_2, \dots)$. This means that the relation \succeq is represented by the utility function U . Here, the space X is treated as a set of consumption outcomes, $(x_0, x_1, x_2, \dots) \in X^\infty$ as a sequence of outcomes consumed over time and the relation \succeq describes the preference of choice of outcomes. We assume that U satisfies the Koopmans recursion $U(x_0, x_1, x_2, \dots) = \varphi(x_0, U(x_1, x_2, \dots))$, where $\varphi: X \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function strictly increasing in its second variable. This is a fundamental axiom in the preference theory in economics (see [1]). Moreover, we assume that \succeq satisfies the property of the preference called "impatience", that is for all $n \geq 1$, $a, b \in X^n$ and for all $(x_0, x_1, \dots) \in X^\infty$, $(a, a, a, \dots) \succeq (b, b, b, \dots) \Leftrightarrow (a, b, x_0, x_1, \dots) \succeq (b, a, x_0, x_1, \dots)$ (see [2]). We consider the Koopmans problem, when the relation \succeq can be represented by another utility function V satisfying the affine recursion $V(x_0, x_1, x_2, \dots) = \alpha(x_0)V(x_1, x_2, \dots) + \beta(x_0)$. We prove that this property holds if and only if there exists a homeomorphic solution F of the system of simultaneous linear functional equations $F(\varphi(x, t)) = \alpha(x)F(t) + \beta(x)$, $x \in X$, $t \in \mathbb{R}$ for some functions $\alpha, \beta: X \rightarrow \mathbb{R}$. If \succeq satisfies impatience, then we give necessary and sufficient conditions that ensure its representation by an affine utility function.

References

- [1] Koopmans, Tjalling C., Peter A. Diamond, and Richard Edmund Williamson. "Stationary utility and time perspective." *Econometrica* 32 (1964): 82-100.
- [2] Bommier, Antoine, Asen Kochov, and François Le Grand. "Ambiguity and endogenous discounting." *J. Math. Econom.* 83 (2019): 48-62.

Weinian Zhang 
Sichuan University, Chengdu, China

Iteration and Equations

Iteration, repeating the same, is a simple operation, but it is important in the informatic era and difficult because of its nonlinearity. This talk surveys the recent advances on iteration and fractional iteration. Furthermore, the advances on various classes of iterative equations are introduced. We also discuss related topics on embeddability of discrete dynamical systems and input-to-state stability of continuous dynamical systems.

Thomas Zürcher

University of Silesia, Katowice, Poland

New solutions to the Matkowski–Wesołowski problem

The Matkowski–Wesołowski problem asks for continuous monotone solutions $\varphi: [0, 1] \rightarrow [0, 1]$ of the equation

$$\varphi(x) = \varphi\left(\frac{x}{2}\right) + \varphi\left(\frac{x+1}{2}\right) - \varphi\left(\frac{1}{2}\right)$$

such that $\varphi(0) = 0$ and $\varphi(1) = 1$.

This talk is based on joint work with Janusz Morawiec, [1]. The goal is to present solutions of this problem that are not Hölder continuous.

References

- [1] Morawiec, Janusz, and Thomas Zürcher. "A new approach with new solutions to the Matkowski and Wesołowski problem." *Aequationes Math.* 95, no. 4 (2021): 761-776.